

**JEE–MAIN EXAMINATION – JUNE, 2022**

**24 June S - 02 Paper Solution**

**SECTION-A**

1. Identify the pair of physical quantities that have same dimensions :

- (A) velocity gradient and decay constant
- (B) wien's constant and Stefan constant
- (C) angular frequency and angular momentum
- (D) wave number and Avogadro number

**Ans. (A)**

**Sol.** Velocity gradient =  $\frac{dV}{dx} = \frac{1}{S}$

$$\lambda = \frac{1}{S}$$

2. The distance between Sun and Earth is R. The duration of year if the distance between Sun and Earth becomes 3R will be :

- (A)  $\sqrt{3}$  years
- (B) 3 years
- (C) 9 years
- (D)  $3\sqrt{3}$  years

**Ans. (D)**

**Sol.**  $T' = T \left( \frac{3R}{R} \right)^{3/2} = 3\sqrt{3} T$

3. A stone of mass m, tied to a string is being whirled in a vertical circle with a uniform speed. The tension in the string is :

- (A) the same throughout the motion
- (B) minimum at the highest position of the circular path
- (C) minimum at the lowest position of the circular path
- (D) minimum when the rope is in the horizontal position

**Ans. (B)**

**Sol.** Theory

4. Two identical charged particles each having a mass 10 g and charge  $2.0 \times 10^{-7}$  C are placed on a horizontal table with a separation of L between them such that they stay in limited equilibrium. If the coefficient of friction between each particle and the table is 0.25, find the value of L. [Use  $g = 10 \text{ ms}^{-2}$ ]

- (A) 12 cm
- (B) 10 cm
- (C) 8 cm
- (D) 5 cm

**Ans. (A)**

**Sol.**  $\frac{kq^2}{L^2} = \mu mg \Rightarrow L = \sqrt{\frac{k}{\mu mg}} q$

5. A Carnot engine takes 5000 kcal of heat from a reservoir at  $727^\circ\text{C}$  and gives heat to a sink at  $127^\circ\text{C}$ . The work done by the engine is :

- (A)  $3 \times 10^6$  J
- (B) Zero
- (C)  $12.6 \times 10^6$  J
- (D)  $8.4 \times 10^6$  J

**Ans. (C)**

**Sol.**  $L = \frac{WD}{Q_H}$

$$\begin{aligned} \Rightarrow WD &= Q_H \left( 1 - \frac{T_L}{T_H} \right) \\ &= 5 \times 10^3 \left( 1 - \frac{400}{1000} \right) \\ &= 3000 \text{ kcal} \end{aligned}$$

6. Two massless springs with spring constants 2 k and k, carry 50 g and 100 g masses at their free ends. These two masses oscillate vertically such that their maximum velocities are equal. Then, the ratio of their respective amplitudes will be :

- (A) 1 : 2
- (B) 3 : 2
- (C) 3 : 1
- (D) 2 : 3

**Ans. (B)**

**Sol.**  $V_{\max} = \omega A$

$$\Rightarrow \frac{A_1}{A_2} = \frac{\omega_2}{\omega_1} = \sqrt{\frac{9}{2} \times \frac{1}{2}} = \frac{3}{2}$$

7. What will be the most suitable combination of three resistors  $A = 2\Omega$ ,  $B = 4\Omega$ ,  $C = 6\Omega$  so that  $\left(\frac{22}{3}\right)\Omega$  is equivalent resistance of combination?

- (A) Parallel combination of A and C connected in series with B.  
 (B) Parallel combination of A and B connected in series with C.  
 (C) Series combination of A and C connected in parallel with B.  
 (D) Series combination of B and C connected in parallel with A.

**Ans. (B)**

**Sol.**  $\Rightarrow \frac{4}{3} + 6 = \frac{22}{3}$

8. The soft-iron is a suitable material for making an electromagnet. This is because soft-iron has :

- (A) low coercivity and high retentivity  
 (B) low coercivity and low permeability  
 (C) high permeability and low retentivity  
 (D) high permeability and high retentivity

**Ans. (C)**

**Sol.** Theory

9. A proton, a deuteron and an  $\alpha$ -particle with same kinetic energy enter into a uniform magnetic field at right angle to magnetic field. The ratio of the radii of their respective circular paths is :

- (A)  $1:\sqrt{2}:\sqrt{2}$                       (B)  $1:1:\sqrt{2}$   
 (C)  $\sqrt{2}:1:1$                               (D)  $1:\sqrt{2}:1$

**Ans. (D)**

**Sol.**  $R = \frac{\sqrt{2km}}{qB} \propto \frac{\sqrt{m}}{q}$

$\frac{\sqrt{m}}{e} : \frac{\sqrt{2m}}{e} : \frac{\sqrt{4m}}{2e}$

$1:\sqrt{2}:1$

10. Given below are two statements :

**Statement-I :** The reactance of an ac circuit is zero. It is possible that the circuit contains a capacitor and an inductor.

**Statement-II :** In ac circuit, the average power delivered by the source never becomes zero.

In the light of the above statements, choose the correct answer from the options given below :

- (A) Both Statement I and Statement II are true.  
 (B) Both Statement I and Statement II are false.  
 (C) Statement I is true but Statement II is false.  
 (D) Statement I is false but Statement II is true.

**Ans. (C)**

**Sol.** if  $R = 0$ ,  $P = 0$

11. Potential energy as a function of  $r$  is given by

$U = \frac{A}{r^{10}} - \frac{B}{r^5}$ , where  $r$  is the interatomic distance,

A and B are positive constants. The equilibrium distance between the two atoms will be :

- (A)  $\left(\frac{A}{B}\right)^{\frac{1}{5}}$                       (B)  $\left(\frac{B}{A}\right)^{\frac{1}{5}}$   
 (C)  $\left(\frac{2A}{B}\right)^{\frac{1}{5}}$                       (D)  $\left(\frac{B}{2A}\right)^{\frac{1}{5}}$

**Ans. (C)**

**Sol.**  $\frac{-10A}{r^{11}} + \frac{5B}{r^6} = 0$


$r^5 = \frac{10A}{5B} = \frac{2A}{B}$

12. An object of mass 5 kg is thrown vertically upwards from the ground. The air resistance produces a constant retarding force of 10 N throughout the motion. The ratio of time of ascent to the time of descent will be equal to : [Use  $g = 10 \text{ ms}^{-2}$ ]

- (A) 1 : 1                                      (B)  $\sqrt{2} : \sqrt{3}$   
 (C)  $\sqrt{3} : \sqrt{2}$                               (D) 2 : 3

**Ans. (B)**

Sol.



$$6t_1^2 = 4t_2^2$$

13. A fly wheel is accelerated uniformly from rest and rotates through 5 rad in the first second. The angle rotated by the fly wheel in the next second, will be :

- (A) 7.5 rad (B) 15 rad  
(C) 20 rad (D) 30 rad

Ans. (B)

Sol.  $5 = \frac{1}{2} \alpha (1)^2$

$$\theta = \frac{1}{2} \alpha (2)^2$$

$$\theta - 5 = 15$$

14. A 100 g of iron nail is hit by a 1.5 kg hammer striking at a velocity of  $60 \text{ ms}^{-1}$ . What will be the rise in the temperature of the nail if one fourth of energy of the hammer goes into heating the nail?

[Specific heat capacity of iron =  $0.42 \text{ Jg}^{-1} \text{ }^\circ\text{C}^{-1}$ ]

- (A)  $675^\circ\text{C}$  (B)  $1600^\circ\text{C}$   
(C)  $160.7^\circ\text{C}$  (D)  $6.75^\circ\text{C}$

Ans. (C)

Sol.  $\frac{1}{2} \times 1.5 \times 60^2 \times \frac{1}{4} = 0.1 \times 420 \times \Delta T$

15. If the charge on a capacitor is increased by 2 C, the energy stored in it increases by 44%. The original charge on the capacitor is (in C) :

- (A) 10 (B) 20  
(C) 30 (D) 40

Ans. (A)

Sol.  $U \propto q^2$   
 $\Rightarrow q_f = 1.2 q$   
 $q_f - q = 2$   
 $\Rightarrow 1.2 q - q = 2$   
 $q = 10$

16. A long cylindrical volume contains a uniformly distributed charge of density  $\rho$ . The radius of cylindrical volume is R. A charge particle (q) revolves around the cylinder in a circular path. The kinetic of the particle is :


- (A)  $\frac{\rho q R^2}{4\epsilon_0}$  (B)  $\frac{\rho q R^2}{2\epsilon_0}$   
(C)  $\frac{q\rho}{4\epsilon_0 R^2}$  (D)  $\frac{4\epsilon_0 R^2}{q\rho}$

Ans. (A)

Sol.  $E = 2\pi r \ell = \frac{\rho \pi r^2 \ell}{\epsilon_0}$

$$qE = \frac{q\rho R^2}{2\epsilon_0 r} = \frac{mv^2}{r}$$

$$mv^2 = \frac{q\rho R^2}{2\epsilon_0}$$



17. An electric bulb is rated as 200 W. What will be the peak magnetic field at 4 m distance produced by the radiations coming from this bulb? Consider this bulb as a point source with 3.5% efficiency.

- (A)  $1.19 \times 10^{-8} \text{ T}$  (B)  $1.71 \times 10^{-8} \text{ T}$   
(C)  $0.84 \times 10^{-8} \text{ T}$  (D)  $3.36 \times 10^{-8} \text{ T}$

Ans. (B)

Sol.  $\frac{\eta P}{4\pi r^2} = \frac{cB_0^2}{2\mu_0}$

$$B_0 = \sqrt{\frac{\mu_0 \eta P}{4\pi c r}}$$

$$\Rightarrow B_0 = \frac{1}{4} \sqrt{\frac{10^{-7} \times 4 \times 3.5}{3 \times 10^8}} = 1.71 \times 10^{-8} \text{ T}$$

18. The light of two different frequencies whose photons have energies 3.8 eV and 1.4 eV respectively, illuminate a metallic surface whose work function is 0.6 eV successively. The ratio of maximum speeds of emitted electrons for the two frequencies respectively will be :

- (A) 1 : 1                      (B) 2 : 1  
(C) 4 : 1                      (D) 1 : 4

**Ans. (B)**

**Sol.**  $\sqrt{\frac{3.8 - 0.6}{1.4 - 0.6}} = \sqrt{\frac{3.2}{0.8}} = 2$

19. Two light beams of intensities in the ratio of 9 : 4 are allowed to interfere. The ratio of the intensity of maxima and minima will be :

- (A) 2 : 3                      (B) 16 : 81  
(C) 25 : 169                (D) 25 : 1

**Ans. (D)**

**Sol.**  $\sqrt{\frac{I_1}{I_2}} = \sqrt{\frac{9}{4}} = \frac{3}{2}$   
 $\left(\frac{\sqrt{I_1} + \sqrt{I_2}}{\sqrt{I_1} - \sqrt{I_2}}\right)^2 = 5^2 = 25$

20. In Bohr's atomic model of hydrogen, let K, P and E are the kinetic energy, potential energy and total energy of the electron respectively. Choose the correct option when the electron undergoes transitions to a higher level :

- (A) All K, P and E increase.  
(B) K decreases. P and E increase.  
(C) P decreases. K and E increase.  
(D) K increases. P and E decrease.



**Ans. (B)**

**Sol.** Based on theory

## SECTION-B

1. A body is projected from the ground at an angle of 45° with the horizontal. Its velocity after 2s is 20 ms<sup>-1</sup>. The maximum height reached by the body during its motion is \_\_\_\_\_m. (use g = 10ms<sup>-2</sup>)

**Ans. (20)**



$v_y = v_x - 20$

$\sqrt{(u_x - 20)^2 + u_x^2} = 20$   
 $\Rightarrow 2u_x^2 - 40u_x = 0$

$\therefore u_x = 20$

2. An antenna is placed in a dielectric medium of dielectric constant 6.25. If the maximum size of that antenna is 5.0 mm. it can radiate a signal of minimum frequency of \_\_\_\_\_GHz.

(Given  $\mu_r = 1$  for dielectric medium)

**Ans. (6)**

**Sol.**  $C' = \frac{C}{\sqrt{\mu_r \epsilon_r}} = \frac{3 \times 10^8}{\sqrt{6.25}} = \frac{3 \times 10^8}{2.5}$

$f\lambda = 1.25 \times 10^8 \text{ s}$


$\Rightarrow f(5 \times 10^{-3} \times 4) = 1.25 \times 10^8$

$f = 6.25 \text{ GHz}$

So  $f \approx 6$

3. A potentiometer wire of length 10 m and resistance  $20 \Omega$  is connected in series with a 25 V battery and an external resistance  $30 \Omega$ . A cell of emf E in secondary circuit is balanced by 250 cm long potentiometer wire. The value of E (in volt) is  $\frac{x}{10}$ . The value of x is \_\_\_\_\_.

Ans. (25)



$$I = \frac{25}{50} = \frac{1}{2} \text{ A}$$

$$\therefore \Delta V = 10 \text{ V}$$

$$10 \text{ m} \rightarrow 10 \text{ V}$$

$$2.5 \text{ m} \rightarrow 2.5 \text{ V}$$

4. Two travelling waves of equal amplitudes and equal frequencies move in opposite directions along a string. They interfere to produce a stationary wave whose equation is given by


$$y = (10 \cos \pi x \sin \frac{2\pi t}{T}) \text{ cm}$$

The amplitude of the particle at  $x = \frac{4}{3}$  cm will be \_\_\_\_\_ cm.

Ans. (5)

Sol.  $10 \cos\left(\frac{4\pi}{3}\right)$

5. In the given circuit- the value of current  $I_L$  will be \_\_\_\_\_ mA.  
(When  $R_L = 1 \text{ k}\Omega$ )



Ans. (5)

Sol.  $I_L = \frac{5}{1000} = 5 \text{ mA}$

6. A sample contains  $10^{-2}$  kg each of two substances A and B with half lives 4 s and 8 s respectively. The ratio of their atomic weights is 1 : 2. The ratio of the amounts of A and B after 16 s is  $\frac{x}{100}$ . The value of x is \_\_\_\_\_.

Ans. (50)

Sol. 
$$N_t = N_0 (0.5)^{\frac{t}{t_{1/2}}}$$

$$= \frac{m}{M} \times N_A (0.5)^{\frac{t}{t_{1/2}}}$$

$$\frac{N_1}{N_2} = \frac{M_2}{M_1} (0.5)^{\left[\frac{1}{T_A} - \frac{1}{T_B}\right] t}$$

$$= 2(0.5)^{16 \times \frac{1}{8}} = \frac{2}{4} = \frac{1}{2} = \frac{x}{100}$$

7. A ray of light is incident at an angle of incidence  $60^\circ$  on the glass slab of refractive index  $\sqrt{3}$ . After refraction, the light ray emerges out from other parallel faces and lateral shift between incident ray and emergent ray is  $4\sqrt{3}$  cm. The thickness of the glass slab is \_\_\_\_\_ cm.

Ans. (12)

**Sol.**  $l = t \sin i \left[ 1 - \frac{\cos i}{\sqrt{\mu^2 - \sin^2 i}} \right]$

$$\Rightarrow 4\sqrt{3} = t \sin 60^\circ \left[ 1 - \frac{\cos 60^\circ}{\sqrt{3 - \frac{3}{4}}} \right]$$

8. A circular coil of 1000 turns each with area  $1\text{m}^2$  is rotated about its vertical diameter at the rate of one revolution per second in a uniform horizontal magnetic field of  $0.07\text{T}$ . The maximum voltage generation will be \_\_\_\_\_ V.

**Ans. (440)**

$$\epsilon_{\text{max}} = BAN\omega$$

**Sol.**

$$\begin{aligned} &= 0.07 \times 1 \times 10^3 \times 2\pi \\ &= 140\pi \approx 440 \end{aligned}$$

9. A monoatomic gas performs a work of  $\frac{Q}{4}$  where Q is the heat supplied to it. The molar heat capacity of the gas will be \_\_\_\_\_ R during this transformation.

Where R is the gas constant.

**Ans. (2)**


**Sol.**  $\Delta Q = \Delta E + WD \Rightarrow Q = \Delta E + \frac{Q}{4}$

$$\Rightarrow n \frac{3R}{2} \Delta T = \Delta E = \frac{3Q}{4}$$

$$\therefore n\Delta T = \frac{Q}{2R}$$

$$\therefore C = 2R$$

10. In an experiment to verify Newton's law of cooling, a graph is plotted between the temperature difference ( $\Delta T$ ) of the water and surroundings and time as shown in figure. The initial temperature of water is taken as  $80^\circ\text{C}$ . The value of  $t_2$  as mentioned in the graph will be \_\_\_\_\_.



**Ans. (16)**

**Sol.**  $T - T_0 = (T_1 - T_0) e^{-\frac{Bt}{ms}}$

$$6\lambda = \ln 1.5$$

$$40 = 60e^{-\lambda(6)} \Rightarrow 6\lambda = \ln 1.5$$

$$20 = 60e^{-\lambda t_2} \Rightarrow t_2 \lambda = \ln 3$$

$$\frac{t_2}{6} = \frac{\ln 3}{\ln 1.5}$$

$$\therefore t_2 = 16.25 \text{ min}$$

$$\text{So } \approx 16$$