







Sol. 1 & 2



6. Adsorption of a gas follows Freundlich adsorption isotherm. If x is the mass of the gas adsorbed on mass m of the adsorbent, the correct plot of $\frac{x}{m}$ versus p is :



Sol. 2


As temp. increases extent of Adsorption decreases
Therefore correct option (2)

$$\frac{x}{m} = Kp^{1/n}$$

$\frac{x}{m}$ v/s $P \rightarrow$ non linear curve

7. The compound that has the largest H-M-H bond angle (M=N, O, S, C) is :
 (1) CH₄ (2) H₂S (3) NH₃ (4) H₂O

Sol. 1



8. The correct statement about probability density (except at infinite distance from nucleus) is :
 (1) It can be zero for 3p orbital (2) It can be zero for 1s orbital
 (3) It can never be zero for 2s orbital (4) It can negative for 2p orbital

Sol. 1

$$\Psi_{R/S}^2 > 0 \text{ always}$$


$$\Psi_{R/S}^2 \text{ can be } = 0; \text{ As '2s' has 1 Radial Node}$$

$$\Psi_R^2 \text{ can never be negative}$$

$$\Psi_R^2 (3P) \text{ can be } = 0 \text{ as 3P has Radial Nodes}$$

Ans. Option (1)

9. The rate constant (k) of a reaction is measured at different temperatures (T), and the data are plotted in the given figure. The activation energy of the reaction in kJ mol⁻¹ is : (R is gas constant)



- Sol. 4 (1) R (2) 2/R (3) 1/R (4) 2R

$$\ln(k) = \ln(A) - \frac{E_a}{R} \left(\frac{1}{T} \right)$$


$$\ln(A) = 10$$

$$\text{Slope} = \frac{-E_a}{R} \times 10^{-3} = -10/5$$

$$E_a = 2000R \text{ J/mol}$$

$$E_a = 2R \text{ KJ/mol}$$


10. The variation of molar conductivity with concentration of an electrolyte (X) in aqueous solution is shown in the given figure.



The electrolyte X is :


- (1) HCl (2) CH₃COOH (3) NaCl (4) KNO₃

Sol. 2




Such type of variation is always for weak electrolyte
Hence Ans (2) CH₃COOH

11. The final major product of the following reaction is :








- (1)
- (2)
- (3)
- (4)


Sol. 3



12. The major product of the following reaction is :



Sol. 3



13. Lattice enthalpy and enthalpy of solution of NaCl are 788 kJ mol^{-1} , and 4 kJ mol^{-1} , respectively. The hydration enthalpy of NaCl is :

(1) -780 kJ mol^{-1}

(2) 784 kJ mol^{-1}

(3) -784 kJ mol^{-1}

(4) 780 kJ mol^{-1}





Sol. 3

$$\Delta H_{\text{sol}} = \text{L.E.} + \Delta H_{\text{hyd}}$$



$$4 = 788 + \Delta H_{\text{Hyd}}$$

$$\Delta H_{\text{Hyd}} = -784 \text{ KJ/mol Ans}$$

14. Reaction of ammonia with excess Cl_2 gives :




Sol. 3



15. Which one of the following polymers is not obtained by condensation polymerisation ?

- (1) Bakelite (2) Nylon 6
(3) Buna-N (4) Nylon 6, 6

Sol. 2



16. Consider the complex ions, trans-[Co(en)₂Cl₂]⁺ (A) and cis-[Co(en)₂Cl₂]⁺ (B)

The correct statement regarding them is :

- (1) Both (A) and (B) can be optically active.
(2) (A) can be optically active, but (B) cannot be optically active.
(3) Both (A) and (B) cannot be optically active.
(4) (A) cannot be optically active, but (B) can be optically active.

Sol. 4

Due to presence of Pos (A) cannot be optically active, but (B) can be optically active

17. An element crystallises in a face-centred cubic (fcc) unit cell with cell edge a. The distance between the centres of two nearest octahedral voids in the crystal lattice is :

- (1) a (2) $\frac{a}{2}$ (3) $\sqrt{2}a$ (4) $\frac{a}{\sqrt{2}}$

Sol. 4

Nearest octahedral voids

One along edge center & other at Body centre

$$\text{Distance} = \sqrt{\left(\frac{a}{2}\right)^2 + \left(\frac{a}{2}\right)^2} = \sqrt{2} \frac{a}{2}$$

$$= \frac{a}{\sqrt{2}} \text{ Ans.}$$

18. The correct order of the ionic radii of O²⁻, N³⁻, F⁻, Mg²⁺, Na⁺ and Al³⁺ is :

- (1) N³⁻ < O²⁻ < F⁻ < Na⁺ < Mg²⁺ < Al³⁺ (2) N³⁻ < F⁻ < O²⁻ < Mg²⁺ < Na⁺ < Al³⁺
(3) Al³⁺ < Na⁺ < Mg²⁺ < O²⁻ < F⁻ < N³⁻ (4) Al³⁺ < Mg²⁺ < Na⁺ < F⁻ < O²⁻ < N³⁻


Sol. 4

all are Isoelectronic

(1) $\frac{\text{N}^{3-}\text{O}^{2-}\text{F}^{-}\text{Na}^{+}\text{Mg}^{2+}\text{Al}^{3+}}{\text{Z} \uparrow, \text{Z}_{\text{eff}} \uparrow, \text{Ionic Radii} \downarrow}$

(2) Al³⁺ < Mg²⁺ < Na⁺ < F⁻ < O²⁻ < N³⁻

19. The increasing order of boiling points of the following compounds is :




(1) I < III < IV < II

(2) IV < I < II < III

(3) I < IV < III < II

(4) III < I < II < IV

Sol. 3



20. The one that is NOT suitable for the removal of permanent hardness of water is :

(1) Ion-exchange method


(2) Calgon's method

(3) Treatment with sodium carbonate

(4) Clark's method

Sol. 4





Clark's method is used for Removal of Temporary hardness



21. For a reaction $X + Y \rightleftharpoons 2Z$, 1.0 mol of X, 1.5 mol of Y and 0.5 mol of Z were taken in a 1 L vessel and allowed to react. At equilibrium, the concentration of Z was 1.0 mol L^{-1} . The equilibrium constant of reaction is _____.

_____ $\frac{x}{15}$. The value of x is _____.

Sol. 16




$$K_{eq} = \frac{(1)^2}{\frac{5}{4} \times \frac{3}{4}} = \frac{16}{15}$$

x = 16 Ans.

22. The volume, in mL, of 0.02 M $K_2Cr_2O_7$ solution required to react with 0.288 g of ferrous oxalate in acidic medium is _____.
(Molar mass of Fe = 56 g mol⁻¹)

Sol. 50 ml




$$\frac{0.02 \times \text{vol} \times 6}{1000} = 3 \times \frac{0.288}{144} \times 100$$

$$\text{Vol.} = \frac{200}{4} = 50 \text{ ml Ans.}$$

23. Considering that $\Delta_0 > P$, the magnetic moment (in BM) of $[Ru(H_2O)_6]^{2+}$ would be _____.

Sol. 0



24. For a dimerization reaction, $2A(g) \rightarrow A_2(g)$ at 298 K, $\Delta U^\ominus = -20 \text{ kJ mol}^{-1}$, $\Delta S^\ominus = -30 \text{ kJ mol}^{-1}$, then the ΔG^\ominus will be _____ J.

Sol. -13538 J



$$\Delta U^\ominus = -20 \text{ kJ}$$

$$\Delta H^\ominus = -20000 + (-1)R \times 298$$

$$\Delta G^\ominus = -20000 - 298R + 30 \times 298$$

$$\Delta G^\ominus = -20,000 + 298 \left(\frac{90 - 25}{3} \right)$$

$$\Delta G^\ominus = 20,000 + \frac{298 \times 65}{3}$$

$$\Delta G^\ominus = -13538 \text{ J}$$

25. The number of chiral carbons present in sucrose is _____.

Sol. 9

