

**MATHEMATICS**  
**JEE-MAIN (July-Attempt) 6 SEPTEMBER**  
**(Shift-1) Paper**

**SECTION - A**

**Q.1** The region represented by  $\{z = x + iy \in \mathbb{C} : |z| - \operatorname{Re}(z) \leq 1\}$  is also given by the inequality:  
 $\{z = x + iy \in \mathbb{C} : |z| - \operatorname{Re}(z) \leq 1\}$

(1)  $y^2 \leq 2\left(x + \frac{1}{2}\right)$     (2)  $y^2 \leq x + \frac{1}{2}$     (3)  $y^2 \geq 2(x + 1)$     (4)  $y^2 \geq x + 1$

**Sol. 1**

$$\{z = x + iy \in \mathbb{C} : |z| - \operatorname{Re}(z) \leq 1\}$$

$$|z| = \sqrt{x^2 + y^2}$$

$$\operatorname{Re}(z) = x$$

$$|z| - \operatorname{Re}(z) \leq 1$$

$$\Rightarrow \sqrt{x^2 + y^2} - x \leq 1$$

$$\Rightarrow \sqrt{x^2 + y^2} \leq 1 + x$$

$$\Rightarrow x^2 + y^2 \leq 1 + x^2 + 2x$$

$$\Rightarrow y^2 \leq 2\left(x + \frac{1}{2}\right)$$

**Q.2** The negation of the Boolean expression  $p \vee (\sim p \wedge q)$  is equivalent to:

(1)  $p \wedge \sim q$     (2)  $\sim p \vee \sim q$     (3)  $\sim p \vee q$     (4)  $\sim p \wedge \sim q$

**Sol. 4**

$$p \vee (\sim p \wedge q)$$

$$(p \wedge \sim p) \wedge (p \vee q)$$

$$t \wedge (p \vee q)$$

$$p \vee q$$

$$\sim (p \vee (\sim p \wedge q)) = \sim (P \vee q)$$

$$= (\sim P) \wedge (\sim q)$$

**Q.3** The general solution of the differential equation  $\sqrt{1+x^2+y^2+x^2y^2} + xy \frac{dy}{dx} = 0$  is:

(where C is a constant of integration)

(1)  $\sqrt{1+y^2} + \sqrt{1+x^2} = \frac{1}{2} \log_e \left( \frac{\sqrt{1+x^2}-1}{\sqrt{1+x^2}+1} \right) + C$

(2)  $\sqrt{1+y^2} - \sqrt{1+x^2} = \frac{1}{2} \log_e \left( \frac{\sqrt{1+x^2}-1}{\sqrt{1+x^2}+1} \right) + C$

(3)  $\sqrt{1+y^2} + \sqrt{1+x^2} = \frac{1}{2} \log_e \left( \frac{\sqrt{1+x^2}+1}{\sqrt{1+x^2}-1} \right) + C$

(4)  $\sqrt{1+y^2} - \sqrt{1+x^2} = \frac{1}{2} \log_e \left( \frac{\sqrt{1+x^2}+1}{\sqrt{1+x^2}-1} \right) + C$

Sol. 3

$$\sqrt{1+x^2+y^2+x^2y^2} + xy \frac{dy}{dx} = 0$$

$$\sqrt{(1+x^2)(1+y^2)} + xy \frac{dy}{dx} = 0$$

$$\frac{\sqrt{(1+x^2)}dx}{x} = -\frac{y}{\sqrt{1+y^2}} dy$$

Integrate the equation

$$\int \frac{\sqrt{1+x^2}}{x} dx = -\int \frac{y}{\sqrt{1+y^2}} dy$$

$$1+x^2 = t^2 \\ 2x dx = 2t dt$$

$$1+y^2 = z^2$$

$$dx = \frac{t}{x} dt$$

$$2y dy = 2z dz$$

$$\int \frac{t \cdot t dt}{t^2 - 1} = -\int \frac{z dz}{z}$$

$$\int \frac{t^2 - 1 + 1}{t^2 - 1} dt = -z + c$$

$$\int 1 dt + \int \frac{1}{t^2 - 1} dt = -z + c$$

$$t + \frac{1}{2} \ln \left( \frac{t-1}{t+1} \right) = -z + c$$

$$\sqrt{1+x^2} + \frac{1}{2} \ln \left( \frac{\sqrt{1+x^2} - 1}{\sqrt{1+x^2} + 1} \right) = -\sqrt{1+y^2} + c$$

$$\sqrt{1+y^2} + \sqrt{1+x^2} = \frac{1}{2} \ln \left( \frac{\sqrt{x^2+1}+1}{\sqrt{x^2+1}-1} \right) + c$$

**Q.4** Let  $L_1$  be a tangent to the parabola  $y^2 = 4(x + 1)$  and  $L_2$  be a tangent to the parabola  $y^2 = 8(x + 2)$  such that  $L_1$  and  $L_2$  intersect at right angles. Then  $L_1$  and  $L_2$  meet on the straight line:

(1)  $x + 2y = 0$

(2)  $x + 2 = 0$

(3)  $2x + 1 = 0$

(4)  $x + 3 = 0$

**Sol. 4**

Let tangent of  $y^2 = 4(x + 1)$

$$L_1 : t_1 y = (x + 1) + t_1^2 \dots\dots(i)$$

and tangent of  $y^2 = 8(x + 2)$

$$L_2 : t_2 y = (x + 2) + 2 t_2^2$$

$$L_1 \perp L_2$$

$$\frac{1}{t_1} \cdot \frac{1}{t_2} = -1$$

$$t_1 t_2 = -1$$

$$t_2(i) - t_1(ii)$$

$$t_1 t_2 y = t_2 (x + 1) + t_2 \cdot t_1^2$$

$$t_1 t_2 y = t_1 (x + 2) + 2 t_2^2 \cdot t_1$$

$$(t_2 - t_1) x + (t_2 - 2t_1) + t_2 t_1 (t_1 - 2t_2) = 0$$

$$(t_2 - t_1) x + (t_2 - 2t_1) - (t_1 - 2t_2) = 0$$

$$(t_2 - t_1) x + 3t_2 - 3t_1 = 0$$

$$\Rightarrow x + 3 = 0$$

**Q.5** The area (in sq. units) of the region  $A = \{(x, y) : |x| + |y| \leq 1, 2y^2 \geq |x|\}$


(1)  $\frac{1}{6}$

(2)  $\frac{5}{6}$

(3)  $\frac{1}{3}$

(4)  $\frac{7}{6}$

**Sol. 2**



$$\text{Total area} = 4 \int_0^{1/2} \left[ (1-x) - \left( \sqrt{\frac{x}{2}} \right) \right] dx$$

$$= 4 \left[ x - \frac{x^2}{2} - \frac{1}{\sqrt{2}} \frac{x^{3/2}}{3/2} \right]_{0}^{1/2}$$

$$= 4 \left[ \frac{1}{2} - \frac{1}{8} - \frac{\sqrt{2}}{3} \left( \frac{1}{2} \right)^{3/2} \right]$$

$$= 4 \times \frac{5}{24} = \frac{5}{6}$$

**Q.6** The shortest distance between the lines  $\frac{x-1}{0} = \frac{y+1}{-1} = \frac{z}{1}$  and  $x + y + z + 1 = 0$ ,  $2x - y + z + 3 = 0$  is:

- (1) 1                      (2)  $\frac{1}{\sqrt{2}}$                       (3)  $\frac{1}{\sqrt{3}}$                       (4)  $\frac{1}{2}$

**Sol. 3**

Plane through line of intersection is  
 $x + y + z + 1 + \lambda(2x - y + z + 3) = 0$   
 It should be parallel to given line  
 $0(1 + 2\lambda) - 1(1 - \lambda) + 1(1 + \lambda) = 0 \Rightarrow \lambda = 0$   
 Plane:  $x + y + z + 1 = 0$   
 Shortest distance of  $(1, -1, 0)$  from this plane

$$= \frac{|1 - 1 + 0 + 1|}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{1}{\sqrt{3}}$$

**Q.7** Let  $a, b, c, d$  and  $p$  be any non zero distinct real numbers such that  $(a^2 + b^2 + c^2)p^2 - 2(ab + bc + cd)p + (b^2 + c^2 + d^2) = 0$ . Then:

- (1)  $a, c, p$  are in G.P.                      (2)  $a, b, c, d$  are in G.P.  
 (3)  $a, b, c, d$  are in A.P.                      (4)  $a, c, p$  are in A.P.

**Sol. 2**

$$(a^2 + b^2 + c^2)p^2 - 2(ab + bc + cd)p + b^2 + c^2 + d^2 = 0$$

$$(a^2p^2 - 2abp + b^2) + [b^2p^2 - 2bcp + c^2] + [c^2p^2 - 2cdp + d^2] = 0$$

$$(ap - b)^2 + (bp - c)^2 + (cp - d)^2 = 0$$

$$ap = b \quad \frac{b}{a} = \frac{c}{b} = \frac{d}{c} = p$$

$$bp = c$$

$$cp = d \quad a, b, c, d \text{ are in G.P.}$$

**Q.8** Two families with three members each and one family with four members are to be seated in a row. In how many ways can they be seated so that the same family members are not separated?

- (1)  $2! 3! 4!$                       (2)  $(3!)^3 \cdot (4!)$                       (3)  $3! (4!)^3$                       (4)  $(3!)^2 \cdot (4!)$

**Sol. 2**

$F_1 \rightarrow 3$  members  
 $F_2 \rightarrow 3$  members  
 $F_3 \rightarrow 4$  members  
 No. of ways can they be seated so that the same family members are not separated  
 $= 3! \times 3! \times 3! \times 4! = (3!)^3 \cdot 4!$

**Q.9** The values of  $\lambda$  and  $\mu$  for which the system of linear equations

$$x + y + z = 2$$

$$x + 2y + 3z = 5$$

$$x + 3y + \lambda z = \mu$$

has infinitely many solutions are, respectively:

(1) 6 and 8

(2) 5 and 8

(3) 5 and 7

(4) 4 and 9

**Sol. 2**

$$x + y + z = 2$$

$$x + 2y + 3z = 5$$

$$x + 3y + \lambda z = \mu$$

has infinitely many solutions

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & \lambda \end{vmatrix} = 0$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$\begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 2 & \lambda - 1 \end{vmatrix} = 0$$

$$(\lambda - 1 - 4) = 0$$

$$\Rightarrow \lambda = 5$$

$$\Delta_3 = \begin{vmatrix} 1 & 1 & 2 \\ 1 & 2 & 5 \\ 1 & 3 & \mu \end{vmatrix} = 0$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$\begin{vmatrix} 1 & 1 & 2 \\ 0 & 1 & 3 \\ 0 & 2 & \mu - 2 \end{vmatrix} = 0$$

$$(\mu - 2) - 6 = 0$$

$$\Rightarrow \mu = 8$$

$$\lambda = 5, \mu = 8$$

**Q.10** Let  $m$  and  $M$  be respectively the minimum and maximum values of

$$\begin{vmatrix} \cos^2 x & 1 + \sin^2 x & \sin 2x \\ 1 + \cos^2 x & \sin^2 x & \sin 2x \\ \cos^2 x & \sin^2 x & 1 + \sin 2x \end{vmatrix}$$

Then the ordered pair  $(m, M)$  is equal to:

- (1)  $(-3, -1)$                       (2)  $(-4, -1)$                       (3)  $(1, 3)$                       (4)  $(-3, 3)$

**Sol. 1**

$$\begin{vmatrix} \cos^2 x & 1 + \sin^2 x & \sin 2x \\ 1 + \cos^2 x & \sin^2 x & \sin 2x \\ \cos^2 x & \sin^2 x & 1 + \sin 2x \end{vmatrix}$$

$$R_1 \rightarrow R_1 - R_2, R_3 \rightarrow R_3 - R_2$$

$$\begin{vmatrix} -1 & 1 & 0 \\ 1 + \cos^2 x & \sin^2 x & \sin 2x \\ -1 & 0 & 1 \end{vmatrix}$$

$$\Rightarrow -1(\sin^2 x) - 1(1 + \cos^2 x + \sin 2x)$$

$$\Rightarrow -\sin^2 x - \cos^2 x - 1 - \sin 2x$$

$$= -2 - \sin 2x$$

$$\therefore \text{minimum value when } \sin 2x = 1$$

$$m = -2 - 1 = -3$$


$$\therefore \text{Maximum value when } \sin 2x = -1$$

$$(m, M) = (-3, -1)$$

**Q.11** A ray of light coming from the point  $(2, 2\sqrt{3})$  is incident at an angle  $30^\circ$  on the line  $x = 1$  at the point A. The ray gets reflected on the line  $x = 1$  and meets x-axis at the point B. Then, the line AB passes through the point:

- (1)  $(4, -\sqrt{3})$                       (2)  $(3, -\frac{1}{\sqrt{3}})$                       (3)  $(3, -\sqrt{3})$                       (4)  $(4, -\frac{\sqrt{3}}{2})$

**Sol. 3**



Equation of P'B  $\rightarrow y - 2\sqrt{3} = \tan 120^\circ (x - 0)$

$$\sqrt{3}x + y = 2\sqrt{3}$$

$(3, -\sqrt{3})$  satisfy the line

**Q.12** Out of 11 consecutive natural numbers if three numbers are selected at random (without repetition), then the probability that they are in A.P. with positive common difference, is:

(1)  $\frac{10}{99}$

(2)  $\frac{5}{33}$

(3)  $\frac{15}{101}$

(4)  $\frac{5}{101}$

**Sol. 2**

**Case-1**

E, O, E, O, E, O, E, O, E, O, E

$$2b = a + c \rightarrow \text{Even}$$

$\Rightarrow$  Both a and c should be either even or odd.

$$P = \frac{{}^6C_2 + {}^5C_2}{{}^{11}C_3} = \frac{5}{33}$$

**Case -2**

O, E, O, E, O, E, O, E, O, E, O

$$P = \frac{{}^5C_2 + {}^6C_2}{{}^{11}C_3} = \frac{5}{33}$$

$$\text{Total probability} = \frac{1}{2} \times \frac{5}{33} + \frac{1}{2} \times \frac{5}{33} = \frac{5}{33}$$

**Q.13** If  $f(x + y) = f(x) f(y)$  and  $\sum_{x=1}^{\infty} f(x) = 2$ ,  $x, y \in \mathbb{N}$ , where  $\mathbb{N}$  is the set of all natural number, then the

value of  $\frac{f(4)}{f(2)}$  is :

(1)  $\frac{2}{3}$

(2)  $\frac{1}{9}$

(3)  $\frac{1}{3}$

(4)  $\frac{4}{9}$

**Sol. 4**

$$f(x + y) = f(x) f(y)$$

\* Put  $x = 1, y = 1$

$$f(2) = (f(1))^2$$

\* Put  $x = 2, y = 1$

$$f(3) = f(2) \cdot f(1) = f((1))^3$$

\* Put  $x = 2, y = 2$

$$f(4) = f((2))^2 = f((1))^4$$

$$f(n) = (f(1))^n$$

$$\sum_{x=1}^{\infty} f(x) = f(1) + f(2) + f(3) + \dots + f(\infty) = 2$$

$$\Rightarrow f(1) + f((1))^2 + f((1))^3 + \dots = 2$$

$$\frac{f(1)}{1 - f(1)} = 2$$

$$f(1) = 2/3$$

$$f(2) = \left(\frac{2}{3}\right)^2, f(4) = \left(\frac{2}{3}\right)^4$$

$$\frac{f(4)}{f(2)} = \frac{(2/3)^4}{(2/3)^2} = \frac{4}{9}$$

**Q.14** If  $\{p\}$  denotes the fractional part of the number  $p$ , then  $\left\{\frac{3^{200}}{8}\right\}$ , is equal to :

(1)  $\frac{5}{8}$

(2)  $\frac{1}{8}$

(3)  $\frac{7}{8}$

(4)  $\frac{3}{8}$

**Sol. 2**

$$\left\{\frac{3^{200}}{8}\right\} = \left\{\frac{9^{100}}{8}\right\} = \left\{\frac{(8+1)^{100}}{8}\right\}$$

$$\left\{\frac{{}^{100}C_0 1^{100} + {}^{100}C_1 (8)1^{99} + {}^{100}C_2 (8^2)1^{98} + \dots + {}^{100}C_{100} 8^{100}}{8}\right\}$$

$$= \left\{\frac{{}^{100}C_0 1^{100} + 8k}{8}\right\}$$

$$= \left\{\frac{1 + 8k}{8}\right\} = \left\{\frac{1}{8} + k\right\} \quad k \in \mathbb{I}$$

$$= \frac{1}{8}$$

**Q.15** Which of the following points lies on the locus of the foot of perpendicular drawn upon any tangent to the ellipse,  $\frac{x^2}{4} + \frac{y^2}{2} = 1$  from any of its foci ?

(1)  $(-1, \sqrt{3})$

(2)  $(-2, \sqrt{3})$


(3)  $(-1, \sqrt{2})$

(4)  $(1, 2)$



**Sol. 4**

Let foot of perpendicular is  $(h,k)$



$$\frac{x^2}{4} + \frac{y^2}{2} = 1 \quad (\text{Given})$$

$$a = 2, b = \sqrt{2}, e = \sqrt{1 - \frac{2}{4}} = \frac{1}{\sqrt{2}}$$

$$\therefore \text{Focus } (ae, 0) = (\sqrt{2}, 0)$$

Equation of tangent

$$y = mx + \sqrt{a^2 m^2 + b^2}$$

$$y = mx + \sqrt{4m^2 + 2}$$

Passes through  $(h,k)$   $(k - mh)^2 = 4m^2 + 2$   
line perpendicular to tangent will have slope

$$-\frac{1}{m}$$

$$y - 0 = -\frac{1}{m}(x - \sqrt{2})$$

$$my = -x + \sqrt{2}$$

$$(h + mk)^2 = 2$$

$$\text{Add equation (1) and (2)} \quad k^2(1 + m^2) + h^2(1 + m^2) = 4(1 + m^2)$$

$$h^2 + k^2 = 4$$

$$x^2 + y^2 = 4 \quad (\text{Auxiliary circle})$$

$\therefore (-1, \sqrt{3})$  lies on the locus.

**Q.16**  $\lim_{x \rightarrow 1} \left( \frac{\int_0^{(x-1)^2} t \cos(t^2) dt}{(x-1) \sin(x-1)} \right)$

(1) is equal to 1      (2) is equal to  $\frac{1}{2}$       (3) does not exist      (4) is equal to  $-\frac{1}{2}$

**Sol**      **Bouns**

$$\lim_{x \rightarrow 1} \left( \frac{\int_0^{(x-1)^2} t \cos(t^2) dt}{(x-1) \sin(x-1)} \right)$$

Using L-Hopital rule

$$= \lim_{x \rightarrow 1} \frac{2(x-1) \cdot (x-1)^2 \cos(x-1)^4 - 0}{(x-1) \cdot \cos(x-1) + \sin(x-1)} \left( \frac{0}{0} \right)$$

$$= \lim_{x \rightarrow 1} \frac{2(x-1)^3 \cdot \cos(x-1)^4}{(x-1) \left[ \cos(x-1) + \frac{\sin(x-1)}{(x-1)} \right]}$$

$$= \lim_{x \rightarrow 1} \frac{2(x-1)^2 \cos(x-1)^4}{(x-1) \left[ \cos(x-1) + \frac{\sin(x-1)}{(x-1)} \right]}$$

$$= \lim_{x \rightarrow 1} \frac{2(x-1)^2 \cos(x-1)^4}{\cos(x-1) + \frac{\sin(x-1)}{(x-1)}}$$

on taking limit

$$= \frac{0}{1+1} = 0$$

**Q.17** If  $\sum_{i=1}^n (x_i - a) = n$  and  $\sum_{i=1}^n (x_i - a)^2 = na$ , ( $n, a > 1$ ) then the standard deviation of  $n$  observations  $x_1, x_2, \dots, x_n$  is :

- (1)  $n\sqrt{a-1}$                       (2)  $\sqrt{na-1}$                       (3)  $a-1$                       (4)  $\sqrt{a-1}$

**Sol. 4**

$$\begin{aligned} \text{S.D.} &= \sqrt{\frac{\sum (x_i - a)^2}{n} - \left(\frac{\sum (x_i - a)}{n}\right)^2} \\ &= \sqrt{\left(\frac{na}{n}\right) - \left(\frac{n}{n}\right)^2} = \sqrt{a-1} \end{aligned}$$

**Q.18** If  $\alpha$  and  $\beta$  be two roots of the equation  $x^2 - 64x + 256 = 0$ . Then the value of

$$\left(\frac{\alpha^3}{\beta^5}\right)^{1/8} + \left(\frac{\beta^3}{\alpha^5}\right)^{1/8} \text{ is :}$$

- (1) 1                      (2) 3                      (3) 2                      (4) 4

**Sol. 3**

$$x^2 - 64x + 256 = 0$$

$$\alpha + \beta = 64$$

$$\alpha\beta = 256$$

$$\begin{aligned} &\left(\frac{\alpha^3}{\beta^5}\right)^{1/8} + \left(\frac{\beta^3}{\alpha^5}\right)^{1/8} \\ &= \frac{\alpha + \beta}{(\alpha\beta)^{5/8}} = \frac{64}{(256)^{5/8}} = \frac{64}{32} = 2 \end{aligned}$$

**Q.19** The position of a moving car at time  $t$  is given by  $f(t) = at^2 + bt + c$ ,  $t > 0$ , where  $a, b$  and  $c$  are real numbers greater than 1. Then the average speed of the car over the time interval  $[t_1, t_2]$  is attained at the point :

- (1)  $(t_1 + t_2)/2$                       (2)  $2a(t_1 + t_2) + b$                       (3)  $(t_2 - t_1)/2$                       (4)  $a(t_2 - t_1) + b$

**Sol. 1**

$$f'(t) = V_{av} = \frac{f(t_2) - f(t_1)}{t_2 - t_1}$$

$$= \frac{a(t_2^2 - t_1^2) + b(t_2 - t_1)}{t_2 - t_1}$$

$$= a(t_1 + t_2) + b = 2at + b$$

$$t = \frac{t_1 + t_2}{2}$$

**Q.20** If  $I_1 = \int_0^1 (1 - x^{50})^{100} dx$  and  $I_2 = \int_0^1 (1 - x^{50})^{101} dx$  such that  $I_2 = \alpha I_1$  then  $\alpha$  equals to :

(1)  $\frac{5050}{5049}$

(2)  $\frac{5050}{5051}$

(3)  $\frac{5051}{5050}$

(4)  $\frac{5049}{5050}$

**Sol. 2**

$$I_1 = \int_0^1 (1 - x^{50})^{100} dx$$

$$I_2 = \int_0^1 (1 - x^{50})(1 - x^{50})^{100} dx$$

$$= \int_0^1 (1 - x^{50})^{100} dx - \int_0^1 x^{50}(1 - x^{50})^{100} dx$$

$$I_2 = I_1 - \int_0^1 \underbrace{x}_{\text{I}} \cdot \underbrace{x^{49}(1 - x^{50})^{100}}_{\text{II}} dx$$

By using by parts  
 $1 - x^{50} = t$

$$\Rightarrow x^{49} dx = \frac{-dt}{50}$$

$$I_2 = I_1 - \left[ x \left( \frac{-1}{50} \right) \frac{(1 - x^{50})^{101}}{101} \right]_0^1 + \int_0^1 \left( \frac{-1}{50} \right) \frac{(1 - x^{50})^{101}}{101} dx$$

$$I_2 = I_1 - 0 + \frac{\int_0^1 (1 - x^{50})^{101} dx}{(-5050)}$$

$$I_2 = I_1 - \frac{I_2}{5050}$$

$$\frac{5051}{5050} I_2 = I_1$$

$$I_2 = \frac{5050}{5051} I_1$$

$$\alpha = \frac{5050}{5051}$$

**Q.21** If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then the greatest value of  $\sqrt{3} |\vec{a} + \vec{b}| + |\vec{a} - \vec{b}|$  is \_\_\_\_\_.

**Sol. 4**

$$\sqrt{3} |\vec{a} + \vec{b}| + |\vec{a} - \vec{b}|$$

$$= \sqrt{3}(\sqrt{2 + 2 \cos \theta}) + \sqrt{2 - 2 \cos \theta}$$


$$= \sqrt{6}(\sqrt{1 + \cos \theta}) + \sqrt{2}(\sqrt{1 - \cos \theta})$$

$$= 2\sqrt{3} \left| \cos \frac{\theta}{2} \right| + 2 \left| \sin \frac{\theta}{2} \right|$$

$$\leq \sqrt{(2\sqrt{3})^2 + (2)^2} = 4$$

**Q.22** Let AD and BC be two vertical poles at A and B respectively on a horizontal ground. If AD = 8 m, BC = 11 m and AB = 10 m; then the distance (in meters) of a point M on AB from the point A such that MD<sup>2</sup> + MC<sup>2</sup> is minimum is \_\_\_\_\_.

**Sol.** 5



$$(MD)^2 = x^2 + 8^2 = x^2 + 64$$

$$(MC)^2 = (10-x)^2 + (11)^2 = (x-10)^2 + 121$$

$$f(x) = (MD)^2 + (MC)^2 = x^2 + 64 + (x-10)^2 + (2)$$

Differentiate

$$f'(x) = 0$$

$$2x + 2(x-10) = 0$$

$$4x = 20 \Rightarrow x = 5$$

$$f''(x) = 4 > 0$$

at  $x = 5$  point of minima

**Q.23** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$f(x) = \begin{cases} x^5 \sin\left(\frac{1}{x}\right) + 5x^2, & x < 0 \\ 0, & x = 0 \\ x^5 \cos\left(\frac{1}{x}\right) + \lambda x^2, & x > 0 \end{cases}$$

The value of  $\lambda$  for which  $f''(0)$  exists, is \_\_\_\_\_.

**Sol. 5**

$$f(x) = \begin{cases} x^5 \sin\left(\frac{1}{x}\right) + 5x^2, & x < 0 \\ 0, & x = 0 \\ x^5 \cos\left(\frac{1}{x}\right) + \lambda x^2, & x > 0 \end{cases}$$


$$f'(x) = \begin{cases} 5x^4 \sin\left(\frac{1}{x}\right) - x^3 \cos\left(\frac{1}{x}\right) + 10x, & x < 0 \\ 0, & x = 0 \\ 5x^4 \cos\left(\frac{1}{x}\right) + x^3 \sin\left(\frac{1}{x}\right) + 2\lambda x, & x > 0 \end{cases}$$

$$f''(x) = \begin{cases} 20x^3 \sin\left(\frac{1}{x}\right) - 5x^2 \cos\left(\frac{1}{x}\right) - 3x^2 \cos\left(\frac{1}{x}\right) - x \sin\left(\frac{1}{x}\right) + 10, & x < 0 \\ 0, & x = 0 \\ 20x^3 \cos\left(\frac{1}{x}\right) + 5x^2 \sin\left(\frac{1}{x}\right) + 3x^2 \sin\left(\frac{1}{x}\right) - x \cos\left(\frac{1}{x}\right) + 2\lambda, & x > 0 \end{cases}$$

$$f''(0^+) = f''(0^-) \\ 2\lambda = 10 \Rightarrow \lambda = 5$$

**Q.24** The angle of elevation of the top of a hill from a point on the horizontal plane passing through the foot of the hill is found to be  $45^\circ$ . After walking a distance of 80 meters towards the top, up a slope inclined at an angle of  $30^\circ$  to the horizontal plane, the angle of elevation of the top of the hill becomes  $75^\circ$ . Then the height of the hill (in meters) is \_\_\_\_\_.

**Sol. 80**



$$x = 80 \cos 30^\circ = 40\sqrt{3}$$

$$y = 80 \sin 30^\circ = 40$$

In  $\triangle ADC$

$$\tan 45^\circ = \frac{h}{x+z} \Rightarrow h = x + z$$

$$\Rightarrow h = 40\sqrt{3} + z \dots (i)$$

In  $\triangle EDF$

$$\tan 75^\circ = \frac{h-y}{z}$$

$$2 + \sqrt{3} = \frac{h-40}{z} \Rightarrow z = \frac{h-40}{2+\sqrt{3}} \dots (ii)$$

Put the value of  $z$  from (i)

$$h - 40\sqrt{3} = \frac{h-40}{2+\sqrt{3}}$$

$$h(1 + \sqrt{3}) = 40(2\sqrt{3} + 3 - 1)$$

$$h(1 + \sqrt{3}) = 80(1 + \sqrt{3})$$

$$h = 80$$

**Q.25** Set A has  $m$  elements and set B has  $n$  elements. If the total number of subsets of A is 112 more than the total number of subsets of B, then the value of  $m \cdot n$  is \_\_\_\_\_.

**Sol. 28**

A & B are set

$$\text{No. of subset of A} = 2^m$$

$$\text{No. of subset of B} = 2^n$$

$$2^m = 2^n + 112$$

$$2^m - 2^n = 112$$

$$2^n(2^{m-n}-1) = 112$$

$$2^n(2^{m-n}-1) = 2^4(2^3-1)$$

$$n = 4 \qquad m - n = 3$$

$$m - 4 = 3 \Rightarrow m = 7$$

$$m \cdot n = 28$$