

CHEMISTRY
JEE-MAIN (February-Attempt) 26
February (Shift-2) Paper


Section - A

1. 2,4-DNP test can be used to identify:






- (1) aldehyde
- (2) halogens
- (3) ether
- (4) amine

Ans. (1)


Sol.



2. Identify A in the following chemical reaction.



Ans. (3)
Sol.



- 3.** The nature of charge on resulting colloidal particles when FeCl₃ is added to excess of hot water is:
- (1) positive
 - (2) neutral
 - (3) sometimes positive and sometimes negative
 - (4) negative


Ans. (1)

Sol. If FeCl₃ is added to excess of hot water, a positively charged sol of hydrated ferric oxide is formed due to adsorption of Fe³⁺ ions.


4. Match **List-I** with **List-II**

List-I


List-II




(i) Wurtz reaction



(ii) Sandmeyer reaction



(iii) Fitting reaction



(iv) Gatterman reaction

Choose the correct answer from the option given below:





(1) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)

(2) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)

(3) (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)

(4) (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)

Ans. (3)



5. In $\overset{1}{\text{C}}\text{H}_2 = \overset{2}{\text{C}} = \overset{3}{\text{C}}\text{H} - \overset{4}{\text{C}}\text{H}_3$ molecule, the hybridization of carbon 1, 2, 3 and 4 respectively are:

- (1) sp^2 , sp , sp^2 , sp^3
- (2) sp^2 , sp^2 , sp^2 , sp^3
- (3) sp^2 , sp^3 , sp^2 , sp^3
- (4) sp^3 , sp , sp^3 , sp^3

Ans. (1)

Sol. $\underset{\text{sp}^2}{\text{C}}\text{H}_2 = \underset{\text{sp}}{\text{C}} = \underset{\text{sp}^2}{\text{C}}\text{H} - \underset{\text{sp}^3}{\text{C}}\text{H}_3$

6. Match List-I with List-II.

List-I

- (a) Sucrose
- (b) Lactose
- (c) Maltose

List-II

- (i) β -D-Galactose and β -D-Glucose
- (ii) α -D-Glucose and β -D-Fructose
- (iii) α -D-Glucose and α -D-Glucose

Choose the correct answer from the options given below:

- (1) (a)-(iii), (b)-(ii), (c)-(i)
- (2) (a)-(iii), (b)-(i), (c)-(ii)
- (3) (a)-(i), (b)-(iii), (c)-(ii)
- (4) (a)-(ii), (b)-(i), (c)-(iii)

Ans. (4)

Sol. Sucrose \rightarrow α -D- Glucose and β -D- Fructose
Lactose \rightarrow β -D- Galactose and β -D- Glucose
Maltose \rightarrow α -D- Glucose and α -D- Glucose

7. Which pair of oxides is acidic in nature?

- (1) N_2O , BaO
- (2) CaO , SiO_2
- (3) B_2O_3 , CaO
- (4) B_2O_3 , SiO_2

Ans. (4)

Sol. B_2O_3 and SiO_2 both are oxides of non-metal and hence are acidic in nature.

8. Calgon is used for water treatment. Which of the following statement is NOT true about calgon?
- (1) Calgon contains the 2nd most abundant element by weight in the earth's crust.
 - (2) It is also known as Graham's salt.
 - (3) It is polymeric compound and is water soluble.
 - (4) It doesnot remove Ca^{2+} ion by precipitation.

Ans. (1)

Sol. $\text{Na}_6(\text{PO}_3)_6$ or $\text{Na}_6\text{P}_6\text{O}_{18}$

Order of abundance of element in earth crust is

$\text{O} > \text{Si} > \text{Al} > \text{Fe} > \text{Ca} > \text{Na} > \text{Mg} > \text{K}$

So second most abundant element in earth crust is Si not Ca.

9. Ceric ammonium nitrate and $\text{CHCl}_3/\text{alc. KOH}$ are used for the identification of functional groups present in _____ and _____ respectively.
- (1) alcohol, amine
 - (2) amine, alcohol
 - (3) alcohol, phenol
 - (4) amine, phenol

Ans. (1)

Sol. Alcohol give positive test with ceric ammonium nitrate and primary amines gives carbyl amine test with $\text{CHCl}_3, \text{KOH}$.

10. Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R.

Assertion A: In TlI_3 , isomorphous to CsI_3 , the metal is present in +1 oxidation state.

Reason R: Tl metals has fourteen *f* electrons in its electronic configuration.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both A and R are correct and R is the correct explanation of A
- (2) A is not correct but R is correct
- (3) Both A and R are correct R is NOT the correct explanation of A
- (4) A is correct but R is not correct

Ans. (3)

Sol. TlI_3 is $\text{Tl}^+ \text{I}_3^-$

CsI_3 is $\text{Cs}^+ \text{I}_3^-$

Thallium shows Tl^+ state due to inert pair effect.


11. The correct order of electron gain enthalpy is:





- (1) $S > Se > Te > O$
- (2) $O > S > Se > Te$
- (3) $S > O > Se > Te$
- (4) $Te > Se > S > O$

Ans. (1)


Sol. Electron gain enthalpy of O is very low due to small size.

12. Identify A in the given chemical reaction.



- (1) 
- (2) 
- (3) 
- (4) 

Ans. (1)



(Internal aldol condensation)

13. Match List-I with List-II

List-I

- (a) Siderite
- (b) Calamine
- (c) Malachite
- (d) Cryolite

List-II

- (i) Cu
- (ii) Ca
- (iii) Fe
- (iv) Al
- (v) Zn



Choose the correct answer from the options given below:

- (1) (a)-(i), (b)-(ii), (c)-(v), (d)-(iii)
- (2) (a)-(iii), (b)-(v), (c)-(i), (d)-(iv)
- (3) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)
- (4) (a)-(iii), (b)-(i), (c)-(v), (d)-(ii)


Ans. (2)

Sol. Siderite - FeCO_3
Calamine - ZnCO_3
Malachite - $\text{CuCO}_3 \cdot \text{Cu(OH)}_2$
Cryolite - Na_3AlF_6

14. Identify A in the given reaction



Ans. (2)



15. Match List-I with List-II.

List-I	List-II
(a) Sodium Carbonate	(i) Deacon
(b) Titanium	(ii) Caster-Kellner
(c) Chlorine	(iii) Van-Arkel
(d) Sodium hydroxide	(iv) Solvay


Choose the correct answer from the option given below:

- (1) (a)-(iii), (b)-(ii), (c)-(i), (d)-(iv)
- (2) (a)-(iv), (b)-(iii), (c)-(i), (d)-(ii)
- (3) (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)
- (4) (a)-(i), (b)-(iii), (c)-(iv), (d)-(ii)


Ans. (2)

Sol. Sodium carbonate Na_2CO_3 & NaHCO_3

Titanium : Van arkel method



Chlorine : Deacon's process



Sodium hydroxide :- Caster-Kellner cell

16. Match List-I with List-II.

List-I (Molecule)	List-II (Bond order)
(a) Ne_2	(i) 1
(b) N_2	(ii) 2
(c) F_2	(iii) 0
(d) O_2	(iv) 3

Choose the correct answer from the options given below:

- (1) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)
- (2) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)
- (3) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)
- (4) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)

Ans. (1)

Sol.

Ne_2O	BO = 0
N_2	BO = 3
F_2	BO = 1
O_2	BO = 2

As per molecular orbital theory

17. Which of the following forms of hydrogen emits low energy β^- particles?

- (1) Proton H^+
- (2) Deuterium 2_1H
- (3) Protium 1_1H
- (4) Tritium 3_1H

Ans. (4)


Sol. Tritium isotope of hydrogen is radioactive and emits low energy β^- particles. It is because of high n/p ratio of tritium which makes nucleus unstable.

18. A. Phenyl methanamine
B. N, N-Dimethylaniline
C. N-Methyl aniline
D. Benzenamine


Choose the correct order of basic nature of the above amines.

- (1) $D > C > B > A$
- (2) $D > B > C > A$
- (3) $A > C > B > D$
- (4) $A > B > C > D$


Ans. (4)



19.




Considering the above reaction, the major product among the following is:



Ans. (3)

Sol.



20. Seliwanoff test and Xanthoproteic test are used for the identification of _____ and _____ respectively

(1) ketoses, proteins

(2) proteins, ketoses

(3) aldoses, ketoses

(4) ketoses, aldoses

Ans. (1)

Sol. Seliwanoff test and Xanthoproteic test are used for identification of 'Ketoses' and proteins respectively.

Section - B

1. The NaNO_3 weighed out to make 50 mL of an aqueous solution containing 70.0 mg Na^+ per mL is _____ g. (Rounded off to the nearest integer)

[Given: Atomic weight in g mol^{-1} . Na: 23; N: 14; O : 16]

Ans. 13

Sol. $\text{Na}^+ = 70 \text{ mg/mL}$

$$\begin{aligned} W_{\text{Na}^+} \text{ in 50mL solution} &= 70 \times 50\text{mg} \\ &= 3500 \text{ mg} \\ &= 3.5 \text{ gm} \end{aligned}$$

$$\text{Moles of } \text{Na}^+ \text{ in 50 ml solution} = \frac{3.5}{23}$$

Moles of $\text{NaNO}_3 = \text{moles of } \text{Na}^+$

$$= \frac{3.5}{23} \text{ mol}$$


$$\text{Mass of } \text{NaNO}_3 = \frac{3.5}{23} \times 85 = 12.934$$

$$\approx 13\text{gm Ans.}$$

2. The number of stereoisomers possible for $[\text{Co}(\text{ox})_2(\text{Br})(\text{NH}_3)]^{2-}$ is _____ [ox = oxalate]

Ans. 3

Sol. $[\text{Co}(\text{ox})_2\text{Br}(\text{NH}_3)]^{2-}$



Total stereoisomer = 2 (OI) + 1 POE (pair of enantiomers) = 3

3. The average S-F bond energy in kJ mol^{-1} of SF_6 is _____. (Rounded off to the nearest integer)

[Given : The values of standard enthalpy of formation of $\text{SF}_6(\text{g})$, $\text{S}(\text{g})$ and $\text{F}(\text{g})$ are - 1100, 275 and 80 kJ mol^{-1} respectively.]

Ans. 309

Sol. $\text{SF}_6(\text{g}) \longrightarrow \text{S}(\text{g}) + 6\text{F}(\text{g})$

$$\Delta H_{\text{reaction}}^{\circ} = 6 \times E_{\text{S-F}} = \Delta H_f^{\circ}[\text{S}(\text{g})] + 6 \times \Delta H_f^{\circ}[\text{F}(\text{g})] - \Delta H_f^{\circ}[\text{SF}_6(\text{g})]$$

$$6 \times E_{\text{S-F}} = 275 + 6 \times 80 - (-1100)$$

$$= 275 + 480 + 1100$$

$$6 \times E_{\text{S-F}} = 1855$$

$$E_{\text{S-F}} = \frac{1855}{6} = 309.1667$$


$\simeq 309 \text{ kJ/mol}$ Ans.

4. Emf of the following cell at 298 K in V is $x \times 10^{-2}$.
 $\text{Zn}|\text{Zn}^{2+} (0.1 \text{ M})||\text{Ag}^+(0.01 \text{ M})|\text{Ag}$
 The value of x is _____. (Rounded off to the nearest integer)

[Given: $E_{\text{Zn}^{2+}/\text{Zn}}^0 = -0.76\text{V}$; $E_{\text{Ag}^+/\text{Ag}}^0 = +0.80\text{V}$; $\frac{2.303RT}{F} = 0.059$]

Ans. 147

Sol. $\text{Zn}(s)|\text{Zn}^{2+}(0.1\text{M})||\text{Ag}^+(0.01\text{M})|\text{Ag}(s)$



$$E^0 = 0.80 + 0.76 = 1.56 ; Q = \left\{ \frac{\text{Zn}^{2+}}{(\text{Ag}^+)^2} \right\}$$

$$E = E^0 - \frac{0.059}{n} \log(Q)$$

$$E = 1.56 - \frac{0.059}{2} \log \left[\frac{0.1}{(0.01)^2} \right]$$

$$E = 1.56 - \frac{0.059}{2} \log \left[(10)^3 \right]$$

$$E = 1.4715 = 147.15 \times 10^{-2} \text{ volt}$$

$$= x \times 10^{-2}$$

$$X = 147.15 \simeq 147 \text{ Ans.}$$

5. A ball weighing 10g is moving with a velocity of 90ms^{-1} . If the uncertainty in its velocity is 5%, then the uncertainty in its position is _____ $\times 10^{-33}\text{m}$. (Rounded off to the nearest integer)

[Given : $h = 6.63 \times 10^{-34} \text{ Js}$]

Ans. 1

Sol. $m = 10 \text{ g} = 10^{-2} \text{ Kg}$

$v = 90 \text{ m/sec.}$

$$\Delta v = v \times 5\% = 90 \times \frac{5}{100} = 4.5 \text{ m / sec}$$

$$m \cdot \Delta v \cdot \Delta x \geq \frac{h}{4\pi}$$

$$10^{-2} \times 4.5 \times \Delta x \geq \frac{6.63 \times 3 \times 10^{-34}}{4 \times \frac{22}{7}}$$


$$\Delta x \geq \frac{6.63 \times 7 \times 2 \times 10^{-34}}{9 \times 4 \times 22 \times 10^{-2}}$$

$$\Delta x \geq 1.17 \times 10^{-33} = x \times 10^{-33}$$

$$x = 1.17 \simeq 1$$

6. In mildly alkaline medium, thiosulphate ion is oxidized by MnO_4^- to "A". The oxidation state of sulphur in "A" is _____.


Ans. 6



∴ Oxidation no. of 'S' = +6 Ans.

7. When 12.2 g of benzoic acid is dissolved in 100g of water, the freezing point of solution was found to be -0.93°C ($K_f(\text{H}_2\text{O}) = 1.86 \text{ K kg mol}^{-1}$). The number (n) of benzoic acid molecules associated (assuming 100% association) is _____.

Ans. 2



$$N = \frac{1}{x} = i \{ \text{As} \quad \alpha = 1 \}$$

$$\Delta T_f = i \times k_f \times m$$

$$0.93 = \frac{1}{n} \times 1.86 \times \frac{12.2 \times 1000}{122 \times 100}$$


$$n = 2$$

8. If the activation energy of a reaction is 80.9 kJ mol^{-1} , the fraction of molecules at 700K, having enough energy to react to form products is e^{-x} . The value of x is _____.

(Rounded off to the nearest integer)

[Use $R = 8.31 \text{ JK}^{-1} \text{ mol}^{-1}$]

Ans. 14




$$\text{Fraction of molecules able to cross energy barrier} = e^{-E_a/RT} = e^{-x}$$

$$x = \frac{E_a}{RT} = \frac{80.9 \times 1000}{8.31 \times 700} = 13.91$$

$$x \simeq 14 \text{ Ans}$$

9. The pH of ammonium phosphate solution, if pK_a of phosphoric acid and pK_b of ammonium hydroxide are 5.23 and 4.75 respectively, is _____.

Ans. 7



$$[H^+] = K_a \times \sqrt{\frac{K_w}{K_a \times K_b}}$$

$$pH = pK_a + \frac{1}{2} \{pK_w - pK_a - pK_b\}$$

$$pH = 5.23 + \frac{1}{2} \{14 - 5.23 - 4.75\}$$

$$pH = 5.23 + \frac{1}{2} (4.02) = 7.24 = 7 \text{ (Nearest integer)}$$

10. The number of octahedral voids per lattice site in a lattice is _____.
(Rounded off to the nearest integer)

Ans. 1

Sol. Assuming FCC

No of lattice sites = 6 face centre + 8 corner = 14

No. of octahedral voids = 13

$$\text{Ratio} = \frac{13}{14} = 0.92857 = 1 \text{ (Nearest integer)}$$