## CARBOXYLIC ACIDS AND IT'S DERIVATIVE, ALIPHATIC AMINES

## EXERCISE # I

- 1. In which reaction product is hydrocarbon?
  - (A) RCOOK  $\xrightarrow{\text{Electrolysis}}$

(B) RCOOAg  $\xrightarrow{l_2/\Delta}$ 

(C)  $CH_3$ – $CH_3$   $\xrightarrow{Cl_2/h\nu}$ 

- (D)  $CH_3 C C1 \xrightarrow{C_2H_5OH}$
- Which of the following set of reaction can not prepare carboxylic acid as the final product: 2.
  - (A)  $R-MgX + O=C=O \xrightarrow{dry \text{ ether}} A \xrightarrow{H_3O^+} (B) R-CN \xrightarrow{(i) SnCl_2 + HCl_3O^+} (B) R-CN \xrightarrow{(ii) H_2O/H^+}$

(C) 
$$CH_2CH_2CH_3 \xrightarrow{KMnO_4,OH} A \xrightarrow{H_3O^+} (D) CH_3(CH_2)_8CH_2OH \xrightarrow{CrO_3} H_2SO_4$$


**3.** In the given reaction,


$$C_{6}H_{5} - C - CH_{3} \xrightarrow{\text{(i) } Br_{2}/KOH} CHBr_{3} + [X]$$


- [X] will be:
- (A)  $C_6H_5$ -CHO
- (B) C<sub>6</sub>H<sub>5</sub>COOH
- (C) C<sub>6</sub>H<sub>5</sub>-CH<sub>2</sub>OH (D) CH<sub>3</sub>COOH


4. Reductive amination of A forms:

$$A: \bigcup_{O}^{NH_2}$$









- 5. In the given reaction:
  - [X] + Acetic anhydride  $\longrightarrow$  Aspirin
  - [X] will be:
  - (A) Benzoic acid

(B) o-methoxybenzoic acid

(C) o-Hydroxybenzoic acid

(D) p-Hydroxybenzoic acid

Arrange following compounds in decreasing order of reactivity for hydrolysis reaction: **6.** 

$$(A) II > IV > I > III$$

(D) 
$$IV > III > II > I$$

7. Which one of the following compounds gives carboxylic acid with HNO<sub>2</sub>?

$$\begin{array}{c} & \text{O} \\ || \\ \text{(A) C}_6\text{H}_5 - \text{C} - \text{Cl} \end{array}$$

$$(C) CH_3 - C - O - C - CH_3$$

8. In the reaction sequence,

$$CH_{3} - C - H \xrightarrow{HCN}_{OH} (A) \xrightarrow{H_{2}O/H^{\oplus}} Product$$

Product will be:

$$(B) \ \ \begin{array}{c} OH \\ | \\ | \\ C-C-CH_3 \\ | \\ H \end{array}$$

(C) Mixture of 
$$CH_3$$
 –  $C$  –  $COOH$  and  $HOOC$  –  $C$  –  $CH_3$  (D)  $CH_3$  –  $CH$  –  $CONH_2$  H

Arrange these esters in decreasing order of ease of esterfication with CH<sub>3</sub>OH/H<sup>⊕</sup>: 9.

(II) 
$$CH_3 - CH - CH_2 - COOH$$
  
 $CH_3$ 

(III) 
$$CH_3 - C - COOH$$
  
 $CH_3$ 

$$\mathrm{(IV)}\,\mathrm{(CH_3-\!CH_2)_3C-\!COOH}$$

(B) 
$$I > II > III > IV$$
 (C)  $III > IV > II > I$ 

(D) 
$$IV > III > II > I$$

# Carboxylic Acids and It's Derivative, Aliphatic Amines

10. Which optically active compound on reduction with LiAlH<sub>4</sub> will give optically inactive compound?

(B) 
$$CH_3 - CH_2 - CH - COOH$$
  
OH

(C) 
$$CH_3 - CH_2 - CH - COOH$$
  
 $CH_2OH$ 

- 11. Which will form lactone on treatment with NaOH?
  - (A) α-Bromo acid
- (B) β-Bromo acid
- (C) β-Hydroxy acid
- (D) δ-Bromo acid

**12.** In the given reaction:


$$\xrightarrow{\text{COOH}} \xrightarrow{\text{HBr}} [X]$$

[X] will be:

13. Correct order of decarboxylation

(a)

(b)



(d)

- (A) a > b > c > d
- (B) c > d > b > a
- (C) c > d > a > b
- (D) d > c > a > b

- **14.** N-Ethyl pthalimide on hydrolysis gives:
  - (A) Methyl alcohol
- (B) Ethyl amine
- (C) Dimethyl amine
- (D) Diethyl amine

### **15.** In the given reaction:

$$[Q] \xleftarrow{\text{NaBH}_4}_{\Delta} \qquad \bigoplus_{O} \xrightarrow{\text{LiAlH}_4} [P]$$

[P] and [Q] respectively be:

(A) 
$$CH_2OH - CH_2 - CH_2 - CH_2OH$$
 and  $OH$ 

(B) on and 
$$CH_2OH - CH_2 - CH_2 - CH - CH_2OH$$
 OH

(C) Both are 
$$O$$
OH

OH (D) Both are 
$$CH_2OH - CH_2 - CH_2 - CH - CH_2OH$$

16. 
$$Ph - C - Cl + O$$

$$MH \longrightarrow A$$
Major 'A' i

(B) 
$$Ph - C - O$$

17. 
$$CH_3NH_2 \xrightarrow{(CH_2)_2O} (A)$$


- (A) 'A' is more basic than  $\mathrm{CH_3NH_2}$
- (B) 'A' is less basic than  $CH_3NH_2$

(C) 'A' is Ter-amine

(D) None

# Carboxylic Acids and It's Derivative, Aliphatic Amines

**18.** Which of the following can released CO<sub>2</sub> with NaHCO<sub>3</sub>.



$$CH_2(COOH)_2$$
  $CH_3$ - $C$ - $COOH$ 

(i)

(ii)

(iii)

- (A) (i), (ii) & (iii)
- (B) (i) & (ii)
- (C) (ii) & (iii)
- (D) (i) & (ii)
- 19. Sodium bicarbonate reacts with salicylic acid to form:

- **20.** Which of the following diazonium salt is relatively stable at 0-5°C:
  - $(A) \ CH_3 N \equiv N \}^{\oplus} Cl^{-}$

(B)  $(CH_3)_2CH-N\equiv N$   $\}^{\oplus}Cl^{-1}$ 

 $(C) C_6H_5-N\equiv N\}^{\oplus}Cl^{-}$ 

(D)  $(CH_3)_3C-N\equiv N$   $\}^{\oplus}Cl^{-1}$ 

- 21. Which is most volatile?
  - (A)  $CH_3CH_2CH_2NH_2$  (B)  $(CH_3)_3N$

- $C_6H_5CONH_2 \xrightarrow{Br_2/OD^{\Theta}} P$ , 'P' is:
  - (A)  $C_6H_5COND_2$

(B)  $C_6H_5ND_2$ 

(C) C<sub>6</sub>H<sub>5</sub>NHD

(D)  $C_6H_5NH_2$ 

In the given reaction: 23.

$$CH_{3} - CH - C - CH_{2} - CH_{3} \xrightarrow{CF_{3}COOOH} [X] \text{ major product}$$

$$CH_{3}$$

[X] will be:

(A) 
$$CH_3 - CH_2 - C - O - CH - CH_3$$
 (B)  $CH_3 - CH - C - O - CH_2 - CH_3$   $CH_3$ 

(B) 
$$CH_3 - CH - C - O - CH_2 - CH_3$$
 $CH_3$ 

$$(C) CH_3 - C - OC(CH_3)_3$$

## **24.** In the given reaction sequence:

$$\begin{array}{c} \operatorname{CH_2-COOH} \\ | \\ \operatorname{CH_2-COOH} \end{array} \xrightarrow{\Delta} (A) \xrightarrow{\operatorname{CH_3-CH_2NH_2/\Delta}} (B)$$

(B) will be:

$$\begin{array}{c} O \\ || \\ || \\ CH_2 - C - NH - C_2H_5 \\ || \\ CH_2 - C - NH - C_2H_5 \\ || \\ O \end{array}$$

$$(B) \begin{tabular}{c} CH_2 - C \\ | \\ CH_2 - C \\ | \\ O \end{tabular} N-C_2H_5$$

(C) 
$$\begin{array}{c} \mathrm{CH}_2\mathrm{-COOH} \\ | & \mathrm{CH}_2\mathrm{-COOH} \end{array}$$

$$(D) \begin{array}{c} O \\ \parallel \\ CH_2 - C - NH - C_2H_5 \\ \mid \\ CH_2 - COOH \end{array}$$


### **25.** In the given reaction :

$$CH_{3}CHO \xrightarrow{(i) \text{ NaCN/HCl}} (A) \xrightarrow{\text{Fenton's}} (B)$$

- (B) will be:
- (A) Acetic acid
- (B) Oxalic acid
- (C) Pyruvic acid
- (D) Citric acid

### EXERCISE # II

1. Which of the following is/are present in mixture of product:



- HO, (A) CH<sub>3</sub>-CH<sub>2</sub>-OH
- Mixture of 1°, 2° and 3° amines can be separated by: 2.
  - (A) Hinsberg's method

(B) Hofmann's isocyanide test

(C) Fractional distillation

(D) NaNO, HCl

- **3.** RCOOR' can be prepared by:
  - (A) Esterification of RCOOH
  - (B) Reaction of CH<sub>3</sub>CH=CH<sub>2</sub> with methanol
  - (C) Baeyer-Villiger oxidation of RCOR' with peroxy acid
  - (D) reaction of RCOCl with R'OH
- Which of the following amine reacts with Hinsberg reagent to give base soluble product:-4.
  - (A) Neopentyl amine
- (B) sec propyl amine (C) diethyl amine
- (D) ethyl methyl amine

5. Which is/are correct reaction(s):

$$(A) \bigcirc -Cl + NH_3 \xrightarrow{\Delta} \bigcirc + NH_4Cl \quad (B) \xrightarrow{\Delta} -Cl + 2NH_3 \xrightarrow{\Delta} NH_2 + NH_4Cl$$

$$(C) \xrightarrow{\hspace{1cm}} Cl + NH_3 \longrightarrow \begin{array}{c} \\ \\ \end{array} + NH_4Cl \\ (D) \xrightarrow{\hspace{1cm}} NH_2 + HNO_2 \xrightarrow{\hspace{1cm}} \begin{array}{c} \\ \\ \\ \end{array} OH \\ \end{array}$$

- Which of the following will form acetyl chloride with PCl<sub>5</sub>? 6.
  - (A) MeCOOH
- (B) MeCOOMe
- (C) MeCOOCOMe
- (D) Me-CONH<sub>2</sub>
- Sodium salt of which compound on electrolysis does not give hydrocarbon: 7.
  - (A)  $C_6H_5COOH$
- (B) HCOOH
- (C) Me<sub>3</sub>C-COOH
- (D) COOH-CH=CH-COOH
- Among the following, which statement is not correct? 8.
  - (A) H<sub>3</sub>C—C—OH will not respond to haloform test
  - (B) Schiff's regent and Schiff's base are different compounds
  - (C) Fehling's solution is a good reagent to detect aromatic aldehydes
  - (D) Both aldehyde and ketone can react with 2,4-dinitrophenylhydrazise reagent

9. 
$$P \xrightarrow{NaOH} P \xrightarrow{CH_3-CH_2-I} R \xrightarrow{H_3O^+} S + T$$

$$N-H \xrightarrow{NaOH} Q + Na_2CO_3$$

If T can evolve efferve scence of  $\mathrm{CO}_2$  with a NaHCO3, then correct statement (s) is/are :

- (A) S & Q can be distinguished by dye azo test
- (B) T is most acidic among all isomeric benzenoid dicarboxylic acid
- (C) Q & S can be distinguished by mustered oil test
- (D) P, Q & T all are soluble in a NaHCO<sub>3</sub>
- **10.** Acetic anhydride and ammonia gives the product:
  - (A) CH<sub>3</sub>CONH<sub>2</sub>
- (B) CH<sub>3</sub>CONHCH<sub>3</sub> (C) CH<sub>3</sub>CN
- (D) CH<sub>3</sub>COONH<sub>4</sub>

11. 
$$CH_3$$
— $C$ — $O$ — $CH_2$ — $CH_3$ — $\frac{HOH/H^{\oplus}}{HOH/H^{\oplus}}$  Ethanoic acid + Ethanol

Isotopic oxygen of water will be present with

- (A) Ethanoic acid
- (B) Ethanol
- (C) After some time it will also be present in some molecules of ester
- (D) None of these
- Which one of the following compounds will give HVZ reaction? **12.**

(A) 
$$\bigcirc$$
 COOH (COOH (D)  $\bigcirc$  COOH

 $RCH_2CONH_2 + NaOBr \longrightarrow ?$ 13.

Rate of reaction will be faster if 'R' is

- $(A) CH_3 -$
- (B)  $C_2H_5$  (C)  $NO_2$  –
- (D) CN -
- Which of the following carboxylic acids do not undergo decarboxylation simply on heating? 14.

(A) 
$$F_3C$$
-C-OH (B)  $H_2C$ =CH-C-OH (C) OH (D)

- **15.** Which of the following compounds will give acetic acid with  $KMnO_4/H^{\oplus}/\Delta$ :
  - (A) CH<sub>3</sub>-CHO

(B) CH<sub>3</sub>-CH=CH-CH<sub>3</sub>

(C)  $CH_3-C\equiv C-CH_3$ 

(D) CH<sub>3</sub>CH<sub>2</sub>OH

# Carboxylic Acids and It's Derivative, Aliphatic Amines

- Hofmann degradation is given by: **16.** 
  - (A) Succinimide
- (B) Acid chloride
- (C) Acid anhydride
- (D) Acetamide
- **17.** The presence of primary amine can be confirmed by its reaction with:
  - (A) HNO<sub>2</sub>
- (B) CHCl<sub>3</sub> + NaOH (C) CS<sub>2</sub> & HgCl<sub>2</sub>
- (D)  $H_2SO_4$
- Total number of compounds which are soluble in hot a NaOH are: **18.** 
  - Salicyclic acid
- (ii) Aspirine
- (iii) Carbolic acid
- (iv) Acetic acid
- (v) Succinic anhydride (vi) Cyclohexanone (vii) Benzene sulphonamide (viii) Cyclohexene

(A) 5

- (B) 6
- (C) 7
- (D) 8
- **19.** Number of oxidation reactions in which organic reactant gets oxidised & one of the major product is carboxylic acid/salt:

(A) 
$$CH_3$$
 NaOI

(B) H OH 
$$H_5IO_6$$
 CHO

$$(D) \bigcirc \stackrel{NC}{\longrightarrow} \stackrel{H_3O^{\oplus}}{\longrightarrow}$$

20. 
$$H_3C-C-O-C=CH_2+H_2O \xrightarrow{H_3O^+} X+Y$$


X and Y are:


(A) X is CH<sub>3</sub>-C


(B) Y is PhCHO

- Which of the following compound react with HNO<sub>2</sub>:









## EXERCISE # III

## **Matching Type Questions**

1. Match the following question:

Column - I

(Reaction)

(A) HOOC 
$$\xrightarrow{\text{CH}_3}$$
 COOH  $\xrightarrow{\Delta}$   $\xrightarrow{\text{COOH}}$   $\xrightarrow{\Delta}$   $C_2H_5$ 

(B) HOOC 
$$\xrightarrow{\text{CH}_3}$$
 COOH  $\xrightarrow{\Delta}$ 

(C) 
$$COOH$$

$$CH_3$$

- COOH COOH

Column-II

(Products)

- (P) Diastereomers
- (Q) Racemic mixture
- (R) Meso comp.
- (S) CO<sub>2</sub> gas will evolve

Match the following question: 2.

Column I

(Organic compounds oxidised by HIO<sub>4</sub>)

- (A) CH<sub>3</sub>COCHO
- (B) 1,2-cyclohexane dione
- (C) PhCH (OH) CHO
- (D) CH<sub>3</sub>CH<sub>2</sub>CH (OH) COCH<sub>3</sub>

Column II

 $(Products\ of\ HIO_{4}\ oxidation)$ 

- (P) PhCH = O + HCOOH
- (Q) CH<sub>3</sub>CH<sub>2</sub>CHO + HOOCCH<sub>3</sub>
- (R) HOOC  $(CH_2)_4COOH$
- (S) CH<sub>3</sub>COOH + HCOOH

# Carboxylic Acids and It's Derivative, Aliphatic Amines

### **3.** Match the following question :

## Column I (Reactions)

(A) 
$$CH_3NH_2 + \bigcirc$$

(B) 
$$(CH_3)_2NH + C_6H_5COCl$$

(C) 
$$\langle \bigcirc \rangle$$
 -CHO +  $\langle \bigcirc \rangle$  -NH<sub>2</sub>

(D) 
$$CH_3NH_2 + \left\langle \bigcirc \right\rangle - SO_2C$$

### **Column II (Products)**

$$(P) \left\langle \bigcirc -N = CH - \bigcirc \right\rangle$$

$$(Q) CH_3 - NH - \bigcup_{O}^{|I|}$$

$$(R) (CH_3)_2N - C \longrightarrow$$

(S) 
$$CH_3NH-C-CH_2CH_2-C-O^2$$

### **4.** Match the following question :

### Column I

### (Correct about product)

(A) 
$$R - CN \xrightarrow{H_3O^+}$$

(B) 
$$R - NH_2 \xrightarrow{CHCl_3} KOH$$

$$(C) \ R-NH_2 \ \tfrac{NaNO_2}{HCl}$$

(D) 
$$R_2NH \xrightarrow{NaNO_2} HCI$$

### Column II

- (P) Product is yellow oily liquid
- (Q) Gives red colour with CAN
- (R) Gives fruity smell with CH<sub>3</sub>OH
- (S) Foul smelling compound is formed.

## 5. Match the following question:

### Column I

(A)  $PhCONH_2 \longrightarrow Ph-C\equiv N$ 

### Column II

 $(P) P_2O_5$ 

(B) 
$$CH_2OH-(CH_2)_2-CHOH-C_6H_5$$
 (Q) LiAlH<sub>4</sub>

(C) 
$$CH_3$$
 – $C$  – $OCH_3$  — $\rightarrow$   $CH_3CH=O$ 

(R)  $H_2/Pd-BaSO_4$ 

(D)  $CH_3COCl \longrightarrow CH_3-CHO$ 

(S) DIBALH

**6.** Match the following question :

### Column I

- (A) RCN \_\_\_\_\_reduction\_\_\_
- (B) RCN  $\frac{(i)CH_3MgBr}{(ii)H_2O}$
- (C) RNC  $\xrightarrow{\text{hydrolysis}}$
- (D)  $RNH_2 \xrightarrow{HNO_2}$

### Column II

- (P) 1° Amine
- (Q) Alcohol
- (R) Ketone
- (S) Acid

7. Find out number of reactions which involve electron difficient nitrogen [Nitrene character] during reaction mechanism.

(a) 
$$Ph$$
  $C = N$   $OH$   $OH$   $OH$ 

(c) 
$$CH_3 - C - NH_2 \xrightarrow{Br_2 + KOH}$$

(e) 
$$CH_3 - C - Cl \xrightarrow{NaN_3 \atop H_2O,\Delta}$$

$$(g) \xrightarrow{H^{\oplus}/H_2O}$$

(b) 
$$H_3C - C - Ph \xrightarrow{PhCO_3H}$$

(d) 
$$Ph - C - OH$$
  $N_3H$ , Dilute  $H_2SO_4$ 

(f) 
$$NH_2 \frac{CHCl_3 + KOH}{}$$

**8.** Of the following amines how many can be seperated by Hoffmann's mustard oil reaction.

$$Ph$$
 $NH_2$ ;  $CH_3$ ;  $NH_2$ ;  $CH_3$ 
 $Ph$ 
 $NH_2$ ;  $Ph$ 
 $NH_2$ ;  $Ph$ 

9. Examine the structure of following compounds, and find out number of compounds that will undergo decarboxylation in presence of heat.

# Carboxylic Acids and It's Derivative, Aliphatic Amines

10. 
$$CH_3CH_2$$
- $CH$ - $COOH \xrightarrow{\Delta} P$  (no. of products)  
 $NH_2$ 

$$\begin{array}{c} CH_{3}CH_{2}\text{--}CH\text{--}COOH & \stackrel{\Delta}{\longrightarrow} & Q \text{ (no. of products)} \\ \downarrow & OH \end{array}$$

$$\begin{array}{c} CH_{3}CH-CH_{2}-COOH \stackrel{\Delta}{\longrightarrow} R \text{ (no. of products)} \\ OH \end{array}$$

$$\begin{array}{c} CH_2\text{--}CH_2\text{--}COOH & \stackrel{\Delta}{\longrightarrow} & S \text{ (no. of products)} \\ OH & \end{array}$$

In all reactions the sum of product is.

## How will you bring about the following transformation:

- 11. Propanoic acid into lactic acid.
- **12.** Ethyl benzene to 2-phenyl propionic acid.
- **13.** Acetamide from acetone.



## **EXERCISE # IV (A) (JEE-MAIN)**

1. Reaction -[AIEEE-2002]

Primary amine + CHCl<sub>3</sub> + KOH → product, here product will be -

- (1) Cyanide
- (2) Isocyanide
- (3) Amine
- (4) Alcohol
- 2. The compound formed in the positive test for nitrogen with the Lassaigne solution of an organic [AIEEE-2004] compound is-

- (1)  $\text{Fe}_{4}[\text{Fe}(\text{CN})_{6}]_{3}$
- (2)  $Na_3[Fe(CN)_6]$  (3)  $Fe(CN)_3$
- $(4) Na_4[Fe(CN)_5)NOS]$
- **3.** Which one of the following methods is neither meant for the synthesis nor for separation of amines-
  - (1) Hofmann method

(2) Hinsberg method

[AIEEE-2005]

(3) Curtius reaction

- (4) Wurtz reaction
- In the chemical reaction,  $CH_2CH_2NH_2 + CHCl_3 + 3KOH \rightarrow (A) + (B) + 3H_2O$ , the compounds (A) 4. [AIEEE-2007] and (B) are respectively -
  - (1) C<sub>2</sub>H<sub>5</sub>CN and 3KCl

(2) CH<sub>3</sub>CH<sub>2</sub> CONH<sub>2</sub> and 3KCl

(3)  $C_2H_5NC$  and  $K_2CO_3$ 

(4) C<sub>2</sub>H<sub>5</sub>NC and 3KCl

**5.** In the chemical reactions, [AIEEE-2010]

$$NH_2$$

$$NaNO_2 \longrightarrow A \longrightarrow B$$
, the compounds 'A' and 'B' respectively are :-


- (1) Nitrobenzene and chlorobenzene
- (2) Nitrobenzene and fluorobenzene
- (3) Phenol and benzene
- (4) Benzene diazonium chloride and fluorobenzene
- **6.** In the chemical reactions

NH,  $\rightarrow$  A  $\xrightarrow{\text{CuCN}}$  B, the compounds A and B respectively are : [AIEEE-2011]

- (1) Fluorobenzene and phenol
- (2) Benzene diazonium chloride and benzonitrile
- (3) Nitrobenzene and chlorobenzene
- (4) Phenol and bromobenzene

# Carboxylic Acids and It's Derivative, Aliphatic Amines

7. Compound (A), C<sub>8</sub>H<sub>9</sub>Br, gives a white precipitate when warmed with alcoholic AgNO<sub>3</sub>. Oxidation of (A) gives an acid (B),  $C_8H_6O_4$ . (B) easily forms anhydride on heating. Identify the compound (A): [AIEEE-2013]



- 8. An organic compound A upon reacting with NH3 gives B. On heating, B gives C. C in presence of KOH reacts with Br<sub>2</sub> to give CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>. A is :-[AIEEE-2013]
  - (1) CH<sub>3</sub>COOH

(2) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>COOH

(3) CH<sub>3</sub>-CH-COOH CH<sub>3</sub>

- (4) CH<sub>3</sub>CH<sub>2</sub>COOH
- 9. On heating an aliphatic primary amine with chloroform & ethenolic potassium hydroside the organic compound formed is [AIEEE-2014]
  - (1) An alkyl cyanide

(2) An alkyl isocyanide

(3) an alkanol

- (4) an alkanediol
- In the reaction  $CH_3COOH \xrightarrow{LiAlH_4} A \xrightarrow{PCl_5} B \xrightarrow{alc.KOH} C$  'C' is [AIEEE-2014] **10.**

- (1) Ethylene
- (2) Acetyl chloride (3) Acetaldehyde
- (4) Acetylene.
- 11. In the presence of a small amount of phosphorous, aliphatic carboxylic acids react with chlorine or bromine to yield a compound in which  $\alpha$  - hydrogen has been replaced by halogen. This reaction is known as: [JEE(Main)-2015]
  - (1) Etard reaction

- (2) Hell Volhard Zelinsky reaction
- (3) Wolff Kischner reaction
- (4) Rosenmund reaction
- **12.** In the Hofmann bromamide degradation reaction, the number of moles of NaOH and Br<sub>2</sub> used per mole of amine produced are: [JEE(Main)-2016]
  - (1) Four moles of NaOH and one mole of Br<sub>2</sub>
  - (2) One mole of NaOH and one mole of Br<sub>2</sub>
  - (3) Four moles of NaOH and two moles of Br<sub>2</sub>
  - (4) Two moles of NaOH and two moles of Br<sub>2</sub>

#### **13.** The major product expected from the following reaction is:

[JEE(Main On-Line)-2017]

$$\begin{array}{c} \text{HO}_2\text{C} & \xrightarrow{\text{CH}_2\text{OH O}} \\ \text{OH} & \xrightarrow{\text{HCl(g)/CCl}_4} \end{array}$$

**14.** The increasing order of basicity of the following compounds is: [JEE(Main)-2018]

(c) 
$$NH_2$$
 (d)  $NHCH_3$ 

$$(1)$$
  $(b) < (a) < (c) < (d)$ 

$$(2)$$
  $(b) < (a) < (d) < (c)$ 

$$(4)$$
  $(a) < (b) < (c) < (d)$ 

Major product of the following reaction is: **15.** 

[JEE Main (Jan)-2019]

$$Cl$$
 $+$ 
 $H_2N$ 
 $O$ 
 $NH_2$ 
 $(2)$  F

(1) Et<sub>3</sub>N (2) Free radical polymerisation

$$(1) \bigcup_{O} \bigcup_{H}^{Cl} \bigcup_{NH_2}^{NH_2}$$

$$(2) \qquad \begin{array}{c} \text{HN} \\ \text{O} \\ \text{NH}_2 \end{array}$$

$$(4) \bigcup_{O}^{CI} \int_{\mathbf{N}} \mathbf{N} \mathbf{N} \mathbf{H}_{2}$$

The major product obtained in the following reaction is: **16.** 

[JEE Main (Jan)-2019]

[JEE Main (Jan)-2019]

$$(3) \qquad \begin{array}{c} O \\ NH \\ CH_3 \end{array}$$

18. Which dicarboxylic acid in presence of a dehydrating agent is least reactive to give an anhydride:

[JEE Main (Jan)-2019]

19. The decreasing order of ease of alkaline hydrolysis for the following esters is:

I

[JEE Main (Jan)-2019]

$$O_2N$$
 COOC<sub>2</sub>H<sub>5</sub>

III

$$(1)$$
 IV  $>$  II  $>$  III  $>$  I

(2) III 
$$>$$
 II  $>$  IV

$$(4) II > III > I > IV$$

**20.** The major product formed in the reaction given below will be :

[**JEE Main (Jan)-2019**]

$$NH_2 \xrightarrow[Aq.HCl,0-5^{\circ}C]{NaNO_2}$$

[JEE Main (Jan)-2019]

$$CH_{3}O \longrightarrow OH \longrightarrow CH_{3} \xrightarrow{\text{(i) dil. HCl/A}} \xrightarrow{\text{(ii) (COOH)}_{2}/} Polymerisation$$

$$(1) \longrightarrow OCOCH_{3} \longrightarrow OCOCH_{3} \longrightarrow OCH_{3} \longrightarrow OCH_{4} \longrightarrow O$$

**22.** An aromatic compound 'A' having molecular formula  $C_7H_6O_2$  on treating with aqueous ammonia and heating forms compound 'B'. The compound 'B' on reaction with molecular bromine and potassium hydroxide provides compound 'C' having molecular formula  $C_6H_7N$ . The structure of 'A' is:

[JEE Main (Jan)-2019]

- 23. A compound 'X' on treatment with Br<sub>2</sub>/NaOH, provided C<sub>3</sub>H<sub>9</sub>N, which gives positive carbylamine test. Compound 'X' is: [JEE Main (Jan)-2019]
  - (1) CH<sub>3</sub>COCH<sub>2</sub>NHCH<sub>3</sub>

(2) CH<sub>3</sub>CH<sub>2</sub>COCH<sub>2</sub>NH<sub>2</sub>

(3) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub>

- (4)  $CH_3CON(CH_3)_2$
- 24. The increasing order of reactivity of the following compounds towards reaction with alkyl halides directly is:

  [JEE Main (Jan)-2019]

$$(A) \qquad (B) \qquad (C) \qquad (D) \qquad NH_2 \qquad (D)$$

(1) (B) < (A) < (D) < (C)

(2) (B) < (A) < (C) < (D)

(3) (A) < (C) < (D) < (B)

(4) (A) < (B) < (C) < (D)

[JEE Main (Jan)-2019]

(1) HO

(2) CH<sub>3</sub> O

(3) CH<sub>3</sub> O

- (4) HO
- **26.** Which of the following amines can be prepared by Gabriel phthalimide reaction?

[JEE Main (Apr)-2019]

- (1) Neo-pentylamine (2) n-butylamine
- (3) triethylamine (4
  - (4) t-butylamine
- **27.** The major product obtained in the following reaction is: [JEE]

Н

[**JEE Main (Apr)-2019**]

$$\begin{array}{c} NH_2 \\ \hline \\ CN & O \end{array}$$


$$(1) \underbrace{ \begin{array}{c} H \\ NCH_3 \\ OH \end{array}}$$


$$(2) \underbrace{ \begin{array}{c} \text{NCH}_3 \\ \text{CN} \end{array} }$$


$$(4) \underbrace{ \begin{pmatrix} H \\ NCHCl_2 \\ CN \end{pmatrix} }_{CN}$$


**28.** The major product of the following reaction is:

[JEE Main (Apr)-2019]









- **29.** Hinsberg's reagent is:
  - (1)  $C_6H_5SO_2Cl$
- (2) C<sub>6</sub>H<sub>5</sub>COCl
- (3) SOCl<sub>2</sub>
- [**JEE Main (Apr)-2019**] (4) (COCl)<sub>2</sub>

[JEE Main (Apr)-2019]

OH (1) CH<sub>3</sub>CHCH<sub>2</sub>CH<sub>2</sub>NHCHO

(2) CH<sub>3</sub>CH=CH-CH<sub>2</sub>NH<sub>2</sub>



OH (4) CH<sub>3</sub>-CH-CH=CH<sub>2</sub>

- (3) CH<sub>3</sub>CHCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>
- 31. Ethylamine (C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub>) can be obtained from N-ethylphthalimide on treatment with :

[JEE Main (Apr)-2019]

- (1) NaBH<sub>4</sub>
- (2) CaH<sub>2</sub>
- (3)  $H_2O$
- (4) NH<sub>2</sub>NH<sub>2</sub>

**32.** The major product 'Y' in the following reaction is:-

[JEE Main (Apr)-2019]

$$Ph \underbrace{\hspace{1cm} CH_3}_{O} \xrightarrow{\hspace{1cm} NaOCl} X \xrightarrow{\hspace{1cm} (i)SOCl_2 \\ \hspace{1cm} (ii)aniline} Y$$

$$(3) \bigvee_{O \to Ph}^{NH_2}$$

$$(4) \qquad \qquad \begin{array}{c} \text{NH}_2 \\ \text{Ph} \end{array}$$

## EXERCISE # IV (B) (JEE ADVANCED)

1. Which of the following carboxylic acids undergo decarboxylation easily:

[IIT 1995]

(A) C<sub>6</sub>H<sub>5</sub>CO-CH<sub>2</sub>COOH

(B) C<sub>6</sub>H<sub>5</sub>COCOOH

(C)  $C_6H_5CH_2$  – COOH OH

- (D)  $C_6H_5CH_2 COOH$  $NH_2$
- 2. The molecular weight of benzoic acid in benzene as determined by depression in freezing point method corresponds to:

  [IIT 1996]
  - (A) Ionization of benzoic acid
  - (B) Dimerisation of benzoic acid
  - (C) Trimerisation of benzoic acid
  - (D) Solvation of benzoic acid
- 3. When propionic acid is treated with aqueous NaHCO<sub>3</sub>, CO<sub>2</sub> is liberated. The 'C' of CO<sub>2</sub> comes from.
  - (A) Methyl group

(B) Carboxylic acid group

(C) methylene group

(D) bicarbonate

[IIT 1999]

**4.** Benzoyl chloride is prepared from benzoic acid by:

[IIT 2000]

- (A)  $Cl_2$ , hv
- (B) SO<sub>2</sub>Cl<sub>2</sub>
- (C) SOCl<sub>2</sub>
- (D) Cl<sub>2</sub>, H<sub>2</sub>O
- **5.** Which of the following acids has the smallest dissociation constant?

[IIT 2002]

(A) CH<sub>3</sub>CHFCOOH

(B) FCH<sub>2</sub>CH<sub>2</sub>COOH

(C) BrCH2CH2COOH

- (D) CH<sub>3</sub>CHBrCOOH
- 6. When benzamide is treated with POCl<sub>3</sub>, the product is:

[IIT 2004]

- (A) Benzonitrile
- (B) Aniline
- (C) Chlorobenzene
- (D) Benzylamine

7. MeO 
$$\longrightarrow$$
 CHO + (X)  $\xrightarrow{\text{CH}_3\text{COONa}}$  MeO  $\longrightarrow$  CH = CH - COCH

The compound (X) is

[IIT 2005]

(A) (CH<sub>3</sub>CO)<sub>2</sub>O

(B) Br  $CH_2$  – COOH

(C) CH<sub>3</sub>COOH

(D) CHO - COOH

**8.** Which of the following reactants on reaction with conc. NaOH followed by acidification gives the following lactone as the only product? [IIT 2006]

- (A) COOCH<sub>3</sub>
- (C) CHO

- (B) COOH
- (D) COOH
- 9. Match the compounds in Column I with their characteristic test(s)/reaction(s) given in Column II. Indicate your answer by darkening the appropriate bubbles of the 4 × 4 matrix given in the ORS. Column-I Column-II [IIT 2008]
  - $(A) \; H_2N \overset{\oplus}{N}\overset{\ominus}{H_3}\overset{\ominus}{C}l$

(P) Sodium fusion extract of the compound gives Prussian blue colour with FeSO<sub>4</sub>

- (C) HO—\(\bigcirc\) \(\time\) \(\tim
- (D)  $O_2N$   $\longrightarrow$   $NH NH_3Br$   $NO_2$

- (Q) Gives positive FeCl<sub>3</sub> test
- (R) Gives white precipitate with  $AgNO_3$
- (S) Reacts with aldehydes to form the

corresponding hydrazone derivative

10. Match each of the compound in Column I with its characteristic reaction(s) in Column II.

## Column-I

- (A) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CN
- (B) CH<sub>3</sub>CH<sub>2</sub>OCOCH<sub>3</sub>
- (C)  $CH_3 CH = CH CH_2OH$
- (D) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>

### Column-II

[IIT 2009]

- (P) Reduction with Pd–C /  $H_2$
- (Q) Reduction with SnCl<sub>2</sub> / HCl
- (R) Development of foul smell on treatment with chloroform and alcoholic KOH
- (S) Reduction with diisobutylaluminium hydride (DIBAL-H)
- (T) Alkaline hydrolysis

[IIT 2011]

$$(B) \bigcirc V - CH_2CI$$


(C) 
$$N$$
O- $CH_2$ 
-Br

$$(D) \bigcirc C$$

$$O - \bigcirc CH_2CI$$

12. With reference the scheme given, which of the given statement(s) about T, U, V & W is/are correct [IIT 2012]

- (A) 'T' is soluble in hot aq NaOH
- (B) 'U' is optically active
- (C) mol formula of W is  $C_{10}H_{18}O_4$
- (D) V gives effervescence with aq NaHCO<sub>3</sub>
- 13. Identify the binary mixtures (s) that can be separated into the individual compounds, by differential extraction, as shown in the given scheme 
  [IIT 2012]



- (A) C<sub>6</sub>H<sub>5</sub>OH and C<sub>6</sub>H<sub>5</sub>COOH
- (B) C<sub>6</sub>H<sub>5</sub>COOH and C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>OH
- (C) C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>OH and C<sub>6</sub>H<sub>5</sub>OH
- (D) C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>OH and C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>COOH

14. The total number of carboxylic acid groups in the product P is

[IIT 2013]

$$\begin{array}{c} O \\ O \\ \hline \\ O \\ \end{array} \begin{array}{c} O \\ \hline \\ \begin{array}{c} 1. \ H_3O^+, \Delta \\ \hline \\ 2. \ O_3 \\ 3. \ H_2O_2 \end{array} \end{array} \right) P$$

15. In the reaction shown below, the major product(s) formed is / are :

[IIT 2014]

$$NH_2 \xrightarrow[CH_2Cl_2]{\text{New acetic anhydride}} product(s)$$

(B) 
$$H_{2}$$
  $CH_{3}$  +  $CH_{3}COOH$ 

(C) 
$$H$$
  $CH_3$   $CH_3 + H_2O$ 

16. Different possible <u>thermal</u> decomposition pathways for peroxyesters are shown below. Match each pathway from List-I with an appropriate structure from List-II and select the correct answer using the code given below the lists.
[IIT 2014]

List-I

List-II

(P) Pathway **P** 

(1) C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub> O CH<sub>3</sub>

(Q) Pathway **Q** 

(2) C<sub>6</sub>H<sub>5</sub> O C<sub>H<sub>3</sub></sub>

(R) Pathway **R** 

(3) C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub> O CH<sub>3</sub> CH<sub>2</sub> CH<sub>2</sub>C<sub>6</sub>H

(S) Pathway S

(4)  $C_6H_5$  O  $CH_3$   $C_2H_5$ 

Code:

P Q R S

- (A) 1 3 4 2
- (B) 2 4 3 1
- (C) 4 1 2 3
- (D) 3 2 1 4

**17.** The major product of the reaction is :

[IIT 2015]

$$(A) \begin{array}{c} H_3C \\ CH_3 \end{array} \begin{array}{c} NH_2 \\ OH \end{array}$$

(B) 
$$H_3C$$
  $CO_2H$   $CH_3$   $OH$ 

(C) 
$$H_3C$$
  $CO_2H$   $CH_3$   $OH$ 

(D) 
$$H_3C$$
  $NH_2$   $CH_3$   $OH$ 

### PARAGRAPH FOR NO. 18 & 19

Treatment of compound  $\mathbf{O}$  with KMnO<sub>4</sub>/H<sup>+</sup> gave  $\mathbf{P}$ , which on heating with ammonia gave The compound  $\mathbf{Q}$  on treatment with Br<sub>2</sub>/NaOH produced  $\mathbf{R}$ . On strong heating,  $\mathbf{Q}$  gave  $\mathbf{S}$ , which on further treatment with ethyl 2-bromopropanoate in the presence of KOH following by acidification, gave a compound  $\mathbf{T}$ .

**18.** The compound  $\mathbf{R}$  is:

- **19.** The compound T is:
  - (A) Glycine
- (B) Alanine
- (C) Valine
- (D) Serine
- 20. The order of basicity among the following compounds is

[IIT-JEE(Adv.)-2017]

(A) II > I > IV > III


(B) IV > II > III > I

(C) I > IV > III > II

(D) IV > I > II > III

### PARAGRAPH FOR NO. 21 & 22

An organic acid  $P(C_{11}H_{12}O_2)$  can easily be oxidized to a dibasic acid which reacts with ethyleneglycol to produce a polymer dacron. Upon ozonolysis, P gives an aliphatic ketone as one of the products. P undergoes the following reaction sequences to furnish R via The compound P also undergoes another set of reactions to produce S. [IIT-JEE(Adv.)-2018]



(There are two questions based on PARAGRAPH "A", the question given below is one of them)

## **21.** The compound $\mathbf{R}$ is

$$(C)$$
  $CO_2H$ 

(D) 
$$CO_2H$$

# **22.** The compound S is

$$(C)$$
  $NH_2$   $(D)$   $NH_2$ 

## **ANSWER-KEY**

### EXERCISE # I

4.

Ans. (C)

- Ans. (A) 2. Ans. (B) **3.**
- 1. Ans. (B)
- 5. **Ans.** (**C**) 6. Ans. (A) 7. Ans. (B) 8. **Ans.** (**C**)
- 9. Ans. (A) **10. Ans.** (C) 11. Ans. (D) **12.** Ans. (B)
- **13. Ans.** (**B**) **14. Ans.** (**B**) **15.** Ans. (A) **16.** Ans. (A)
- **17.** Ans. (B) 18. **Ans.** (**C**) **19.** Ans. (B) **20.** Ans. (C)
- 21. **Ans.** (**B**) 22. Ans. (B) 23. Ans. (A) 24. Ans. (B)
- 25. **Ans.** (C)

### **EXERCISE # II**

- Ans. (A,C,D)2. **Ans.** (**A**,**C**) 3. Ans. (A,C,D)4. **Ans.** (**A**,**B**) 1.
- 5. Ans. (A,C,D)7. 6. Ans. (A,B,C)**Ans.** (**B**,**C**) 8. Ans. (C)
- 9. Ans. (A,B,D)Ans. (A) 11. Ans. (A,C) **12. Ans.** (**B**,**C**) **10.**
- 13. Ans. (B) **14.** Ans. (A,B,C)**15.** Ans. (A,B,C,D)**16. Ans.** (**A**,**D**)
- **17.** Ans. (A,B,C)Ans. (B) **19.** Ans. (A,B,D)**20. Ans.** (**C,D**) **18.**
- 21. Ans. (B,C,D)

### **EXERCISE** # III

- 1. Ans. (A) $\rightarrow$ P, S; (B) $\rightarrow$ Q, S; (C) $\rightarrow$ P, S; (D) $\rightarrow$ R
- 2. Ans. (A) $\rightarrow$ S; (B) $\rightarrow$ R; (C) $\rightarrow$ P; (D) $\rightarrow$ Q 3. Ans. (A) $\rightarrow$ S; (B) $\rightarrow$ R; (C) $\rightarrow$ P; (D) $\rightarrow$ Q
- 4. Ans. (A) $\rightarrow$ R; (B) $\rightarrow$ S; (C) $\rightarrow$ Q; (D) $\rightarrow$ P 5. Ans. (A) $\rightarrow$ P; (B) $\rightarrow$ Q; (C) $\rightarrow$ S; (D) $\rightarrow$ R, S
- 6. Ans. (A) $\rightarrow$ P; (B) $\rightarrow$ R; (C) $\rightarrow$ P, S; (D) $\rightarrow$ Q
- 7. **10.** Ans. (9) Ans. (4) 8. **Ans.** (4) Ans. (5)

## EXERCISE # IV (A) (JEE-MAIN)

- 1. Ans. (2) 2. Ans. (1) 3. Ans. (4) 4. Ans. (4)
- 6. 5. Ans. (4) Ans. (2) 7. Ans. (4) 8. Ans. (4)
- 9. Ans. (2) 10. **Ans.** (1) 11. Ans. (2) **12.** Ans. (1)
- Ans. (1) 13.
- **14.** Ans. (2)
- Sol. Order of base nature depends on electron donation tendency.

In compound NH nitrogen is sp<sup>2</sup> hybridized so least basic among all given compound.

compound — is very strong nitrogeneous organic base as lone pair of one nitrogen delocalize in

resonance and make another nitrogen negativly charged and conjugate acid have two equivalent resonating structure.

Thus it is most basic in given compouds.

NHCH<sub>3</sub> (secondary amine) more basic than NH<sub>2</sub> (primary amine)

### 15. Ans. (4)

Sol. 
$$Cl$$
  $Cl$   $NH_2$   $Cl$   $NH_2$   $Cl$   $NH$   $O$   $O$ 

NH<sub>2</sub>(a) will wact as nucleophile as (b) is having delocalised lonepair.

$$\begin{array}{c|c} Cl & NH_2 & \hline \\ O & Polymerisation \\ \hline \end{array}$$

### 16. Ans. (3)

Sol. 
$$CH_3 - C - O - C - CH_3$$

$$OH$$

$$OH$$

$$OH$$

$$OH$$

$$NH-C-CH_3$$

### 17. Ans. (3)

Sol. 
$$O$$

$$C$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{2}-CH_{3}$$

$$CH_{3}-CH_{3}$$

$$CH_{3}-CH_{3}$$

$$CH_{3}-CH_{3}$$

$$CH_{3}-CH_{3}$$

18. Ans. (4)

Sol. Adipic acid 
$$CO_2H$$
– $(CH_2)_4$ – $CO_2H$   $\xrightarrow{\text{dehydrating}}$  7 membered cyclic anhydride (Very unstable)

19. Ans. (2)

**Sol.** More is the electrophilic character of carbonyl group of ester faster is the alkaline hydrolysis.

20. Ans. (Bonus)

21. Ans. (2)

22. Ans. (3)

#### 23. Ans. (3)

Sol. 
$$[X] \xrightarrow{Br_2} C_3H_9N \xrightarrow{CHCl_3} CH_3CH_2CH_2-NC$$

Hoff mann's Carbylamine

Bromaide Reaction

degradation

Thus [X] must be aride with oen carbon more than is amine.

Thus [X] is CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub>

#### 24. Ans. (2)

Sol. Nucleophilicity order

$$\bigcup_{0}^{O} NH < \bigcup_{0}^{O} NH_{2} < \bigcup_{0}^{CN} NH_{2} < \bigcup_{0}^{N} NH_{2}$$

$$A \qquad C \qquad D$$

### 25.

Sol.

$$H_3C$$
O
O
O
O
O

#### **26.** Ans. (2)

Gabriel phthalimide synthesis:

$$\begin{array}{c} O \\ O \\ O \\ O \\ O \\ \end{array}$$
 NH 
$$\begin{array}{c} 1.KOH \\ \hline 2.R-X \\ (1^{\circ}halide\ S_{N}) \end{array}$$
 O 
$$O \\ O \\ H_{3}O^{+} \\ \end{array}$$
 COOH 
$$R-NH_{2}+ \begin{array}{c} COOH \\ \hline \end{array}$$

### 27. Ans. (1)

[Benzene Sulphonyl chloride]

30. Ans. (1)

as NH<sub>2</sub> is a better nucleophile than OH.

31. Ans. (4)

Sol. 
$$\bigcirc N-\text{Et} \xrightarrow{H_2N-NH_2} C_2H_5NH_2$$

reagent is  $NH_2$ - $NH_2$  byproduct will be

32. Ans. (1)

Sol. Ph 
$$CH_3$$
 NaOCl Ph  $C$  NH  $C$  Ph  $C$  NH  $C$  Ph

## EXERCISE # IV (B) (JEE ADVANCED)

1. Ans. (A)

2. Ans. (B)

3. Ans. (D)

4. Ans. (C)

5. Ans. (C)

6. Ans. (A)

7. Ans. (A)

8. Ans. (C)

9. Ans. (A) $\rightarrow$ R, S; (B) $\rightarrow$ P, Q; (C) $\rightarrow$ P, Q, R; (D) $\rightarrow$ P,S

10. Ans. (A) $\rightarrow$ P, Q, S, T; (B) $\rightarrow$ P, S, T; (C) $\rightarrow$ P; (D) $\rightarrow$ R

11. Ans. (A)

12. Ans. (A,C,D)

13. Ans. (B,D)

14. Ans. (2)

15. Ans. (A)

16. Ans. (A)

17. Ans. (C)

18. Ans. (A)

19. Ans. (B)

**Solution 18 & 19.** 

Q to R is Hoffmann's bromamide degradation reaction

S to T is Gabriel's phthalimide sysnthesis

20. Ans. (D) IV > I > II > III

Sol. Basic strength  $\infty$  stability of conjugated acid.

$$\propto$$
 + M / +H / +I

21. Ans. (A)

22. Ans. (B)

## **Solution 21 & 22.**