



## SOLUTIONS TO JEE (ADVANCED) - 2015

(PAPER - 02)

Time : 3 Hours

Maximum Marks : 240

### READ THE INSTRUCTIONS CAREFULLY

#### QUESTION PAPER FORMAT AND MARKING SCHEME :

1. The question paper has three parts: Physics, Chemistry and Mathematics. Each part has three sections.
2. Section 1 contains 8 questions. The answer to each question is a single digit integer ranging from 0 to 9 (both inclusive).  
**Marking Scheme:** +4 for correct answer and 0 in all other cases.
3. Section 2 contains 8 multiple choice questions with one or more than one correct option.  
**Marking Scheme:** +4 for correct answer, 0 if not attempted and -2 in all other cases.
4. Section 3 contains 2 “paragraph” type questions. Each paragraph describes an experiment, a situation or a problem. Two multiple choice questions will be asked based on this paragraph. One or more than one option can be correct.  
**Marking Scheme:** +4 for correct answer, 0 if not attempted and -2 in all other cases.


## PART-I: PHYSICS

### Section 1 (Maximum Marks: 32)

- This section contains **EIGHT** questions.
- The answer to each question is a **SINGLE DIGIT INTEGER** ranging from 0 to 9, **both** inclusive.
- For each question, darken the bubble corresponding to the correct integer in the ORS.
- Marking scheme:
  - +4 If the bubble corresponding to the answer is darkened.
  - 0 In all other cases.

1. An electron in an excited state of  $\text{Li}^{2+}$  ion has angular momentum  $3h/2\pi$ . The de Broglie wavelength of the electron in this state is  $p\pi a_0$  (where  $a_0$  is the Bohr radius). The value of  $p$  is

\*2. A large spherical mass  $M$  is fixed at one position and two identical point masses  $m$  are kept on a line passing through the centre of  $M$  (see figure). The point masses are connected by a rigid massless rod of length  $\ell$  and this assembly is free to move along the line connecting them. All three masses interact only through their mutual gravitational interaction. When the point mass nearer to  $M$  is at a distance  $r = 3\ell$  from  $M$ , the tension in the rod is zero for  $m = k\left(\frac{M}{288}\right)$ . The value of  $k$  is




3. The energy of a system as a function of time  $t$  is given as  $E(t) = A^2 \exp(-\alpha t)$ , where  $\alpha = 0.2 \text{ s}^{-1}$ . The measurement of  $A$  has an error of 1.25%. If the error in the measurement of time is 1.50%, the percentage error in the value of  $E(t)$  at  $t = 5 \text{ s}$  is

\*4. The densities of two solid spheres A and B of the same radii  $R$  vary with radial distance  $r$  as  $\rho_A(r) = k\left(\frac{r}{R}\right)$  and  $\rho_B(r) = k\left(\frac{r}{R}\right)^5$ , respectively, where  $k$  is a constant. The moments of inertia of the individual spheres about axes passing through their centres are  $I_A$  and  $I_B$ , respectively. If  $\frac{I_B}{I_A} = \frac{n}{10}$ , the value of  $n$  is


\*5. Four harmonic waves of equal frequencies and equal intensities  $I_0$  have phase angles  $0, \pi/3, 2\pi/3$  and  $\pi$ . When they are superposed, the intensity of the resulting wave is  $nI_0$ . The value of  $n$  is

6. For a radioactive material, its activity  $A$  and rate of change of its activity  $R$  are defined as  $A = -\frac{dN}{dt}$  and  $R = -\frac{dA}{dt}$ , where  $N(t)$  is the number of nuclei at time  $t$ . Two radioactive sources P (mean life  $\tau$ ) and Q (mean life  $2\tau$ ) have the same activity at  $t = 0$ . Their rates of change of activities at  $t = 2\tau$  are  $R_P$  and  $R_Q$ , respectively. If  $\frac{R_P}{R_Q} = \frac{n}{e}$ , then the value of  $n$  is

7. A monochromatic beam of light is incident at  $60^\circ$  on one face of an equilateral prism of refractive index  $n$  and emerges from the opposite face making an angle  $\theta(n)$  with the normal (see the figure). For  $n = \sqrt{3}$  the value of  $\theta$  is  $60^\circ$  and  $\frac{d\theta}{dn} = m$ . The value of  $m$  is



8. In the following circuit, the current through the resistor  $R (=2\Omega)$  is  $I$  Amperes. The value of  $I$  is




### Section 2 (Maximum Marks: 32)

- This section contains **EIGHT** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct.
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- Marking scheme:
  - +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened.
  - 0 If none of the bubbles is darkened
  - 2 In all other cases

9. A fission reaction is given by  ${}_{92}^{236}\text{U} \rightarrow {}_{54}^{140}\text{Xe} + {}_{38}^{94}\text{Sr} + x + y$ , where  $x$  and  $y$  are two particles. Considering  ${}_{92}^{236}\text{U}$  to be at rest, the kinetic energies of the products are denoted by  $K_{\text{Xe}}$ ,  $K_{\text{Sr}}$ ,  $K_x(2\text{MeV})$  and  $K_y(2\text{MeV})$ , respectively. Let the binding energies per nucleon of  ${}_{92}^{236}\text{U}$ ,  ${}_{54}^{140}\text{Xe}$  and  ${}_{38}^{94}\text{Sr}$  be 7.5 MeV, 8.5 MeV and 8.5 MeV respectively. Considering different conservation laws, the correct option(s) is(are)
- (A)  $x = n$ ,  $y = n$ ,  $K_{\text{Sr}} = 129\text{MeV}$ ,  $K_{\text{Xe}} = 86\text{MeV}$   
 (B)  $x = p$ ,  $y = e^-$ ,  $K_{\text{Sr}} = 129\text{MeV}$ ,  $K_{\text{Xe}} = 86\text{MeV}$   
 (C)  $x = p$ ,  $y = n$ ,  $K_{\text{Sr}} = 129\text{MeV}$ ,  $K_{\text{Xe}} = 86\text{MeV}$   
 (D)  $x = n$ ,  $y = n$ ,  $K_{\text{Sr}} = 86\text{MeV}$ ,  $K_{\text{Xe}} = 129\text{MeV}$

- \*10. Two spheres P and Q of equal radii have densities  $\rho_1$  and  $\rho_2$ , respectively. The spheres are connected by a massless string and placed in liquids  $L_1$  and  $L_2$  of densities  $\sigma_1$  and  $\sigma_2$  and viscosities  $\eta_1$  and  $\eta_2$ , respectively. They float in equilibrium with the sphere P in  $L_1$  and sphere Q in  $L_2$  and the string being taut (see figure). If sphere P alone in  $L_2$  has terminal velocity  $\vec{V}_p$  and Q alone in  $L_1$  has terminal velocity  $\vec{V}_q$ , then




- (A)  $\frac{|\vec{V}_p|}{|\vec{V}_q|} = \frac{\eta_1}{\eta_2}$  (B)  $\frac{|\vec{V}_p|}{|\vec{V}_q|} = \frac{\eta_2}{\eta_1}$   
 (C)  $\vec{V}_p \cdot \vec{V}_q > 0$  (D)  $\vec{V}_p \cdot \vec{V}_q < 0$

11. In terms of potential difference  $V$ , electric current  $I$ , permittivity  $\epsilon_0$ , permeability  $\mu_0$  and speed of light  $c$ , the dimensionally correct equation(s) is(are)


- (A)  $\mu_0 I^2 = \epsilon_0 V^2$  (B)  $\epsilon_0 I = \mu_0 V$   
 (C)  $I = \epsilon_0 c V$  (D)  $\mu_0 c I = \epsilon_0 V$

12. Consider a uniform spherical charge distribution of radius  $R_1$  centred at the origin  $O$ . In this distribution, a spherical cavity of radius  $R_2$ , centred at  $P$  with distance  $OP = a = R_1 - R_2$  (see figure) is made. If the electric field inside the cavity at position  $\vec{r}$  is  $\vec{E}(\vec{r})$ , then the correct statement(s) is(are)



- (A)  $\vec{E}$  is uniform, its magnitude is independent of  $R_2$  but its direction depends on  $\vec{r}$   
 (B)  $\vec{E}$  is uniform, its magnitude depends on  $R_2$  and its direction depends on  $\vec{r}$   
 (C)  $\vec{E}$  is uniform, its magnitude is independent of  $a$  but its direction depends on  $\vec{a}$   
 (D)  $\vec{E}$  is uniform and both its magnitude and direction depend on  $\vec{a}$

- \*13. In plotting stress versus strain curves for two materials P and Q, a student by mistake puts strain on the y-axis and stress on the x-axis as shown in the figure. Then the correct statement(s) is(are)




- (A) P has more tensile strength than Q  
 (B) P is more ductile than Q  
 (C) P is more brittle than Q  
 (D) The Young's modulus of P is more than that of Q

- \*14. A spherical body of radius  $R$  consists of a fluid of constant density and is in equilibrium under its own gravity. If  $P(r)$  is the pressure at  $r$  ( $r < R$ ), then the correct option(s) is(are)


- (A)  $P(r = 0) = 0$  (B)  $\frac{P(r = 3R/4)}{P(r = 2R/3)} = \frac{63}{80}$   
 (C)  $\frac{P(r = 3R/5)}{P(r = 2R/5)} = \frac{16}{21}$  (D)  $\frac{P(r = R/2)}{P(r = R/3)} = \frac{20}{27}$

15. A parallel plate capacitor having plates of area  $S$  and plate separation  $d$ , has capacitance  $C_1$  in air. When two dielectrics of different relative permittivities ( $\epsilon_1 = 2$  and  $\epsilon_2 = 4$ ) are introduced between the two plates as shown in the figure, the capacitance becomes  $C_2$ . The ratio  $\frac{C_2}{C_1}$  is



- (A)  $6/5$  (B)  $5/3$   
 (C)  $7/5$  (D)  $7/3$

- \*16. An ideal monoatomic gas is confined in a horizontal cylinder by a spring loaded piston (as shown in the figure). Initially the gas is at temperature  $T_1$ , pressure  $P_1$  and volume  $V_1$  and the spring is in its relaxed state. The gas is then heated very slowly to temperature  $T_2$ , pressure  $P_2$  and volume  $V_2$ . During this process the piston moves out by a distance  $x$ . Ignoring the friction between the piston and the cylinder, the correct statement(s) is(are)




- (A) If  $V_2 = 2V_1$  and  $T_2 = 3T_1$ , then the energy stored in the spring is  $\frac{1}{4}P_1V_1$   
 (B) If  $V_2 = 2V_1$  and  $T_2 = 3T_1$ , then the change in internal energy is  $3P_1V_1$   
 (C) If  $V_2 = 3V_1$  and  $T_2 = 4T_1$ , then the work done by the gas is  $\frac{7}{3}P_1V_1$   
 (D) If  $V_2 = 3V_1$  and  $T_2 = 4T_1$ , then the heat supplied to the gas is  $\frac{17}{6}P_1V_1$

### SECTION 3 (Maximum Marks: 16)

- This section contains **TWO** paragraphs
- Based on each paragraph, there will be **TWO** questions
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS
- Marking scheme:
  - +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened
  - 0 If none of the bubbles is darkened
  - 2 In all other cases

### PARAGRAPH 1


Light guidance in an optical fiber can be understood by considering a structure comprising of thin solid glass cylinder of refractive index  $n_1$  surrounded by a medium of lower refractive index  $n_2$ . The light guidance in the structure takes place due to successive total internal reflections at the interface of the media  $n_1$  and  $n_2$  as shown in the figure. All rays with the angle of incidence  $i$  less than a particular value  $i_m$  are confined in the medium of refractive index  $n_1$ . The numerical aperture (NA) of the structure is defined as  $\sin i_m$ .



17. For two structures namely  $S_1$  with  $n_1 = \sqrt{45}/4$  and  $n_2 = 3/2$ , and  $S_2$  with  $n_1 = 8/5$  and  $n_2 = 7/5$  and taking the refractive index of water to be  $4/3$  and that of air to be 1, the correct option(s) is(are)
- (A) NA of  $S_1$  immersed in water is the same as that of  $S_2$  immersed in a liquid of refractive index  $\frac{16}{3\sqrt{15}}$
- (B) NA of  $S_1$  immersed in liquid of refractive index  $\frac{6}{\sqrt{15}}$  is the same as that of  $S_2$  immersed in water
- (C) NA of  $S_1$  placed in air is the same as that of  $S_2$  immersed in liquid of refractive index  $\frac{4}{\sqrt{15}}$ .
- (D) NA of  $S_1$  placed in air is the same as that of  $S_2$  placed in water
18. If two structures of same cross-sectional area, but different numerical apertures  $NA_1$  and  $NA_2$  ( $NA_2 < NA_1$ ) are joined longitudinally, the numerical aperture of the combined structure is
- (A)  $\frac{NA_1 NA_2}{NA_1 + NA_2}$  (B)  $NA_1 + NA_2$
- (C)  $NA_1$  (D)  $NA_2$

### PARAGRAPH 2

In a thin rectangular metallic strip a constant current  $I$  flows along the positive  $x$ -direction, as shown in the figure. The length, width and thickness of the strip are  $\ell$ ,  $w$  and  $d$ , respectively. A uniform magnetic field  $\vec{B}$  is applied on the strip along the positive  $y$ -direction. Due to this, the charge carriers experience a net deflection along the  $z$ -direction. This results in accumulation of charge carriers on the surface PQRS and appearance of equal and opposite charges on the face opposite to PQRS. A potential difference along the  $z$ -direction is thus developed. Charge accumulation continues until the magnetic force is balanced by the electric force. The current is assumed to be uniformly distributed on the cross section of the strip and carried by electrons.



19. Consider two different metallic strips (1 and 2) of the same material. Their lengths are the same, widths are  $w_1$  and  $w_2$  and thicknesses are  $d_1$  and  $d_2$ , respectively. Two points K and M are symmetrically located on the opposite faces parallel to the  $x$ - $y$  plane (see figure).  $V_1$  and  $V_2$  are the potential differences between K and M in strips 1 and 2, respectively. Then, for a given current  $I$  flowing through them in a given magnetic field strength  $B$ , the correct statement(s) is(are)
- (A) If  $w_1 = w_2$  and  $d_1 = 2d_2$ , then  $V_2 = 2V_1$       (B) If  $w_1 = w_2$  and  $d_1 = 2d_2$ , then  $V_2 = V_1$   
 (C) If  $w_1 = 2w_2$  and  $d_1 = d_2$ , then  $V_2 = 2V_1$       (D) If  $w_1 = 2w_2$  and  $d_1 = d_2$ , then  $V_2 = V_1$
20. Consider two different metallic strips (1 and 2) of same dimensions (lengths  $\ell$ , width  $w$  and thickness  $d$ ) with carrier densities  $n_1$  and  $n_2$ , respectively. Strip 1 is placed in magnetic field  $B_1$  and strip 2 is placed in magnetic field  $B_2$ , both along positive  $y$ -directions. Then  $V_1$  and  $V_2$  are the potential differences developed between K and M in strips 1 and 2, respectively. Assuming that the current  $I$  is the same for both the strips, the correct option(s) is(are)
- (A) If  $B_1 = B_2$  and  $n_1 = 2n_2$ , then  $V_2 = 2V_1$       (B) If  $B_1 = B_2$  and  $n_1 = 2n_2$ , then  $V_2 = V_1$   
 (C) If  $B_1 = 2B_2$  and  $n_1 = n_2$ , then  $V_2 = 0.5V_1$       (D) If  $B_1 = 2B_2$  and  $n_1 = n_2$ , then  $V_2 = V_1$

JEE


## PART-II: CHEMISTRY

### SECTION 1 (Maximum Marks: 32)


- This section contains **EIGHT** questions
- The answer to each question is a **SINGLE DIGIT INTEGER** ranging from 0 to 9, both inclusive
- For each question, darken the bubble corresponding to the correct integer in the ORS
- Marking scheme:  
 +4 If the bubble corresponding to the answer is darkened  
 0 In all other cases

\*21. In dilute aqueous  $\text{H}_2\text{SO}_4$ , the complex diaquodioxalatoferrate(II) is oxidized by  $\text{MnO}_4^-$ . For this reaction, the ratio of the rate of change of  $[\text{H}^+]$  to the rate of change of  $[\text{MnO}_4^-]$  is

\*22. The number of hydroxyl group(s) in Q is



23. Among the following, the number of reaction(s) that produce(s) benzaldehyde is



24. In the complex acetyl bromidodicarbonylbis(triethylphosphine)iron(II), the number of Fe–C bond(s) is

25. Among the complex ions,  $[\text{Co}(\text{NH}_2\text{-CH}_2\text{-CH}_2\text{-NH}_2)_2\text{Cl}_2]^+$ ,  $[\text{CrCl}_2(\text{C}_2\text{O}_4)_2]^{3-}$ ,  $[\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2]^+$ ,  $[\text{Fe}(\text{NH}_3)_2(\text{CN})_4]^-$ ,  $[\text{Co}(\text{NH}_2\text{-CH}_2\text{-CH}_2\text{-NH}_2)_2(\text{NH}_3)\text{Cl}]^{2+}$  and  $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]^{2+}$ , the number of complex ion(s) that show(s) *cis-trans* isomerism is

\*26. Three moles of  $\text{B}_2\text{H}_6$  are completely reacted with methanol. The number of moles of boron containing product formed is

27. The molar conductivity of a solution of a weak acid HX (0.01 M) is 10 times smaller than the molar conductivity of a solution of a weak acid HY (0.10 M). If  $\lambda_{\text{X}^-}^0 \approx \lambda_{\text{Y}^-}^0$ , the difference in their  $\text{pK}_a$  values,  $\text{pK}_a(\text{HX}) - \text{pK}_a(\text{HY})$ , is (consider degree of ionization of both acids to be  $\ll 1$ )




28. A closed vessel with rigid walls contains 1 mol of  $^{238}_{92}\text{U}$  and 1 mol of air at 298 K. Considering complete decay of  $^{238}_{92}\text{U}$  to  $^{206}_{82}\text{Pb}$ , the ratio of the final pressure to the initial pressure of the system at 298 K is


**SECTION 2 (Maximum Marks: 32)**

- This section contains **EIGHT** questions
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS
- Marking scheme:  
 +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened  
 0 If none of the bubbles is darkened  
 -2 In all other cases






- \*29. One mole of a monoatomic real gas satisfies the equation  $p(V - b) = RT$  where  $b$  is a constant. The relationship of interatomic potential  $V(r)$  and interatomic distance  $r$  for the gas is given by








30. In the following reactions, the product **S** is



31. The major product **U** in the following reactions is



32. In the following reactions, the major product **W** is



\*33. The correct statement(s) regarding, (i) HClO, (ii) HClO<sub>2</sub>, (iii) HClO<sub>3</sub> and (iv) HClO<sub>4</sub>, is (are)

- (A) The number of Cl = O bonds in (ii) and (iii) together is two  
 (B) The number of lone pairs of electrons on Cl in (ii) and (iii) together is three  
 (C) The hybridization of Cl in (iv) is sp<sup>3</sup>  
 (D) Amongst (i) to (iv), the strongest acid is (i)

34. The pair(s) of ions where BOTH the ions are precipitated upon passing  $\text{H}_2\text{S}$  gas in presence of dilute  $\text{HCl}$ , is(are)
- (A)  $\text{Ba}^{2+}$ ,  $\text{Zn}^{2+}$  (B)  $\text{Bi}^{3+}$ ,  $\text{Fe}^{3+}$   
 (C)  $\text{Cu}^{2+}$ ,  $\text{Pb}^{2+}$  (D)  $\text{Hg}^{2+}$ ,  $\text{Bi}^{3+}$
- \*35. Under hydrolytic conditions, the compounds used for preparation of linear polymer and for chain termination, respectively, are
- (A)  $\text{CH}_3\text{SiCl}_3$  and  $\text{Si}(\text{CH}_3)_4$  (B)  $(\text{CH}_3)_2\text{SiCl}_2$  and  $(\text{CH}_3)_3\text{SiCl}$   
 (C)  $(\text{CH}_3)_2\text{SiCl}_2$  and  $\text{CH}_3\text{SiCl}_3$  (D)  $\text{SiCl}_4$  and  $(\text{CH}_3)_3\text{SiCl}$
36. When  $\text{O}_2$  is adsorbed on a metallic surface, electron transfer occurs from the metal to  $\text{O}_2$ . The **TRUE** statement(s) regarding this adsorption is(are)
- (A)  $\text{O}_2$  is physisorbed (B) heat is released  
 (C) occupancy of  $\pi_{2p}^*$  of  $\text{O}_2$  is increased (D) bond length of  $\text{O}_2$  is increased

### SECTION 3 (Maximum Marks: 16)

- This section contains **TWO** paragraphs
- Based on each paragraph, there will be **TWO** questions
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS
- Marking scheme:
  - +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened
  - 0 In none of the bubbles is darkened
  - 2 In all other cases

### PARAGRAPH 1


When 100 mL of 1.0 M  $\text{HCl}$  was mixed with 100 mL of 1.0 M  $\text{NaOH}$  in an insulated beaker at constant pressure, a temperature increase of  $5.7^\circ\text{C}$  was measured for the beaker and its contents (**Expt. 1**). Because the enthalpy of neutralization of a strong acid with a strong base is a constant ( $-57.0 \text{ kJ mol}^{-1}$ ), this experiment could be used to measure the calorimeter constant. In a second experiment (**Expt. 2**), 100 mL of 2.0 M acetic acid ( $K_a = 2.0 \times 10^{-5}$ ) was mixed with 100 mL of 1.0 M  $\text{NaOH}$  (under identical conditions to **Expt. 1**) where a temperature rise of  $5.6^\circ\text{C}$  was measured.

(Consider heat capacity of all solutions as  $4.2 \text{ J g}^{-1} \text{ K}^{-1}$  and density of all solutions as  $1.0 \text{ g mL}^{-1}$ )





- \*37. Enthalpy of dissociation (in  $\text{kJ mol}^{-1}$ ) of acetic acid obtained from the **Expt. 2** is
- (A) 1.0 (B) 10.0  
 (C) 24.5 (D) 51.4
- \*38. The pH of the solution after **Expt. 2** is
- (A) 2.8 (B) 4.7  
 (C) 5.0 (D) 7.0

### PARAGRAPH 2





In the following reactions



39. Compound X is



40. The major compound Y is



  
JEE

## PART-III: MATHEMATICS

### Section 1 (Maximum Marks: 32)

- This section contains **EIGHT** questions.
- The answer to each question is a **SINGLE DIGIT INTEGER** ranging from 0 to 9, both inclusive.
- For each question, darken the bubble corresponding to **the** correct integer in the ORS.
- Marking scheme:
  - +4 If the bubble corresponding to the answer is darkened.
  - 0 In all other cases.

41. Suppose that  $\vec{p}, \vec{q}$  and  $\vec{r}$  are three non-coplanar vectors in  $\mathbb{R}^3$ . Let the components of a vector  $\vec{s}$  along  $\vec{p}, \vec{q}$  and  $\vec{r}$  be 4, 3 and 5, respectively. If the components of this vector  $\vec{s}$  along  $(-\vec{p} + \vec{q} + \vec{r}), (\vec{p} - \vec{q} + \vec{r})$  and  $(-\vec{p} - \vec{q} + \vec{r})$  are  $x, y$  and  $z$ , respectively, then the value of  $2x + y + z$  is

\*42. For any integer  $k$ , let  $\alpha_k = \cos\left(\frac{k\pi}{7}\right) + i \sin\left(\frac{k\pi}{7}\right)$ , where  $i = \sqrt{-1}$ . The value of the expression

$$\frac{\sum_{k=1}^{12} |\alpha_{k+1} - \alpha_k|}{\sum_{k=1}^3 |\alpha_{4k-1} - \alpha_{4k-2}|}$$
 is

\*43. Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first eleven terms is 6 : 11 and the seventh term lies in between 130 and 140, then the common difference of this A.P. is

\*44. The coefficient of  $x^9$  in the expansion of  $(1+x)(1+x^2)(1+x^3) \dots (1+x^{100})$  is

\*45. Suppose that the foci of the ellipse  $\frac{x^2}{9} + \frac{y^2}{5} = 1$  are  $(f_1, 0)$  and  $(f_2, 0)$  where  $f_1 > 0$  and  $f_2 < 0$ . Let  $P_1$  and  $P_2$  be two parabolas with a common vertex at  $(0, 0)$  and with foci at  $(f_1, 0)$  and  $(2f_2, 0)$ , respectively. Let  $T_1$  be a tangent to  $P_1$  which passes through  $(2f_2, 0)$  and  $T_2$  be a tangent to  $P_2$  which passes through  $(f_1, 0)$ . The  $m_1$  is the slope of  $T_1$  and  $m_2$  is the slope of  $T_2$ , then the value of  $\left(\frac{1}{m_1^2} + m_2^2\right)$  is

46. Let  $m$  and  $n$  be two positive integers greater than 1. If

$$\lim_{\alpha \rightarrow 0} \left( \frac{e^{\cos(\alpha^n)} - e}{\alpha^m} \right) = -\left(\frac{e}{2}\right)$$

then the value of  $\frac{m}{n}$  is

47. If

$$\alpha = \int_0^1 (e^{9x+3\tan^{-1}x}) \left( \frac{12+9x^2}{1+x^2} \right) dx$$

where  $\tan^{-1}x$  takes only principal values, then the value of  $\left(\log_e |1+\alpha| - \frac{3\pi}{4}\right)$  is

48. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous odd function, which vanishes exactly at one point and  $f(1) = \frac{1}{2}$ . Suppose that  $F(x) = \int_{-1}^x f(t) dt$  for all  $x \in [-1, 2]$  and  $G(x) = \int_{-1}^x t|f(f(t))| dt$  for all  $x \in [-1, 2]$ . If  $\lim_{x \rightarrow 1} \frac{F(x)}{G(x)} = \frac{1}{14}$ , then the value of  $f\left(\frac{1}{2}\right)$  is

**Section 2 (Maximum Marks: 32)**

- This section contains **EIGHT** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct.
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- Marking scheme:
  - +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened.
  - 0 If none of the bubbles is darkened
  - 2 In all other cases

49. Let  $f'(x) = \frac{192x^3}{2 + \sin^4 \pi x}$  for all  $x \in \mathbb{R}$  with  $f\left(\frac{1}{2}\right) = 0$ . If  $m \leq \int_{1/2}^1 f(x) dx \leq M$ , then the possible values of  $m$  and  $M$  are
- (A)  $m = 13, M = 24$  (B)  $m = \frac{1}{4}, M = \frac{1}{2}$   
 (C)  $m = -11, M = 0$  (D)  $m = 1, M = 12$
- \*50. Let  $S$  be the set of all non-zero real numbers  $\alpha$  such that the quadratic equation  $\alpha x^2 - x + \alpha = 0$  has two distinct real roots  $x_1$  and  $x_2$  satisfying the inequality  $|x_1 - x_2| < 1$ . Which of the following intervals is(are) a subset(s) of  $S$ ?
- (A)  $\left(-\frac{1}{2}, -\frac{1}{\sqrt{5}}\right)$  (B)  $\left(-\frac{1}{\sqrt{5}}, 0\right)$   
 (C)  $\left(0, \frac{1}{\sqrt{5}}\right)$  (D)  $\left(\frac{1}{\sqrt{5}}, \frac{1}{2}\right)$
- \*51. If  $\alpha = 3 \sin^{-1}\left(\frac{6}{11}\right)$  and  $\beta = 3 \cos^{-1}\left(\frac{4}{9}\right)$ , where the inverse trigonometric functions take only the principal values, then the correct option(s) is(are)
- (A)  $\cos \beta > 0$  (B)  $\sin \beta < 0$   
 (C)  $\cos(\alpha + \beta) > 0$  (D)  $\cos \alpha < 0$
- \*52. Let  $E_1$  and  $E_2$  be two ellipses whose centers are at the origin. The major axes of  $E_1$  and  $E_2$  lie along the x-axis and the y-axis, respectively. Let  $S$  be the circle  $x^2 + (y - 1)^2 = 2$ . The straight line  $x + y = 3$  touches the curves  $S, E_1$  and  $E_2$  at  $P, Q$  and  $R$ , respectively. Suppose that  $PQ = PR = \frac{2\sqrt{2}}{3}$ . If  $e_1$  and  $e_2$  are the eccentricities of  $E_1$  and  $E_2$ , respectively, then the correct expression(s) is(are)
- (A)  $e_1^2 + e_2^2 = \frac{43}{40}$  (B)  $e_1 e_2 = \frac{\sqrt{7}}{2\sqrt{10}}$   
 (C)  $|e_1^2 - e_2^2| = \frac{5}{8}$  (D)  $e_1 e_2 = \frac{\sqrt{3}}{4}$

\*53. Consider the hyperbola  $H : x^2 - y^2 = 1$  and a circle  $S$  with center  $N(x_2, 0)$ . Suppose that  $H$  and  $S$  touch each other at a point  $P(x_1, y_1)$  with  $x_1 > 1$  and  $y_1 > 0$ . The common tangent to  $H$  and  $S$  at  $P$  intersects the  $x$ -axis at point  $M$ . If  $(l, m)$  is the centroid of the triangle  $\Delta PMN$ , then the correct expression(s) is(are)

(A)  $\frac{dl}{dx_1} = 1 - \frac{1}{3x_1^2}$  for  $x_1 > 1$

(B)  $\frac{dm}{dx_1} = \frac{x_1}{3(\sqrt{x_1^2 - 1})}$  for  $x_1 > 1$

(C)  $\frac{dl}{dx_1} = 1 + \frac{1}{3x_1^2}$  for  $x_1 > 1$

(D)  $\frac{dm}{dy_1} = \frac{1}{3}$  for  $y_1 > 0$

54. The option(s) with the values of  $a$  and  $L$  that satisfy the following equation is(are)

$$\frac{\int_0^{4\pi} e^t (\sin^6 at + \cos^4 at) dt}{\int_0^{\pi} e^t (\sin^6 at + \cos^4 at) dt} = L ?$$

(A)  $a = 2, L = \frac{e^{4\pi} - 1}{e^{\pi} - 1}$

(B)  $a = 2, L = \frac{e^{4\pi} + 1}{e^{\pi} + 1}$

(C)  $a = 4, L = \frac{e^{4\pi} - 1}{e^{\pi} - 1}$

(D)  $a = 4, L = \frac{e^{4\pi} + 1}{e^{\pi} + 1}$

55. Let  $f, g : [-1, 2] \rightarrow \mathbb{R}$  be continuous functions which are twice differentiable on the interval  $(-1, 2)$ . Let the values of  $f$  and  $g$  at the points  $-1, 0$  and  $2$  be as given in the following table:

	$x = -1$	$x = 0$	$x = 2$
$f(x)$	3	6	0
$g(x)$	0	1	-1

In each of the intervals  $(-1, 0)$  and  $(0, 2)$  the function  $(f - 3g)''$  never vanishes. Then the correct statement(s) is(are)

(A)  $f'(x) - 3g'(x) = 0$  has exactly three solutions in  $(-1, 0) \cup (0, 2)$

(B)  $f'(x) - 3g'(x) = 0$  has exactly one solution in  $(-1, 0)$

(C)  $f'(x) - 3g'(x) = 0$  has exactly one solution in  $(0, 2)$

(D)  $f'(x) - 3g'(x) = 0$  has exactly two solutions in  $(-1, 0)$  and exactly two solutions in  $(0, 2)$

56. Let  $f(x) = 7\tan^8 x + 7\tan^6 x - 3\tan^4 x - 3\tan^2 x$  for all  $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ . Then the correct expression(s) is(are)

(A)  $\int_0^{\pi/4} xf'(x) dx = \frac{1}{12}$

(B)  $\int_0^{\pi/4} f(x) dx = 0$

(C)  $\int_0^{\pi/4} xf'(x) dx = \frac{1}{6}$

(D)  $\int_0^{\pi/4} f(x) dx = 1$

### SECTION 3 (Maximum Marks: 16)

- This section contains **TWO** paragraphs.
- Based on each paragraph, there will be **TWO** questions
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct
- For each question, darken the bubble(s) corresponding to all the correct option(s) in the ORS.
- Marking scheme:
  - +4 If only the bubble(s) corresponding to all the correct option(s) is(are) darkened.
  - 0 If none of the bubbles is darkened
  - 2 In all other cases

#### PARAGRAPH 1

Let  $F : \mathbb{R} \rightarrow \mathbb{R}$  be a thrice differentiable function. Suppose that  $F(1) = 0$ ,  $F(3) = -4$  and  $F'(x) < 0$  for all  $x \in (1/2, 3)$ . Let  $f(x) = xF(x)$  for all  $x \in \mathbb{R}$ .

57. The correct statement(s) is(are)
- (A)  $f'(1) < 0$  (B)  $f(2) < 0$   
(C)  $f'(x) \neq 0$  for any  $x \in (1, 3)$  (D)  $f'(x) = 0$  for some  $x \in (1, 3)$
58. If  $\int_1^3 x^2 F'(x) dx = -12$  and  $\int_1^3 x^3 F''(x) dx = 40$ , then the correct expression(s) is(are)
- (A)  $9f'(3) + f'(1) - 32 = 0$  (B)  $\int_1^3 f(x) dx = 12$   
(C)  $9f'(3) - f'(1) + 32 = 0$  (D)  $\int_1^3 f(x) dx = -12$

#### PARAGRAPH 2

Let  $n_1$  and  $n_2$  be the number of red and black balls, respectively, in box I. Let  $n_3$  and  $n_4$  be the number of red and black balls, respectively, in box II.

59. One of the two boxes, box I and box II, was selected at random and a ball was drawn randomly out of this box. The ball was found to be red. If the probability that this red ball was drawn from box II is  $\frac{1}{3}$ , then the correct option(s) with the possible values of  $n_1, n_2, n_3$  and  $n_4$  is(are)
- (A)  $n_1 = 3, n_2 = 3, n_3 = 5, n_4 = 15$  (B)  $n_1 = 3, n_2 = 6, n_3 = 10, n_4 = 50$   
(C)  $n_1 = 8, n_2 = 6, n_3 = 5, n_4 = 20$  (D)  $n_1 = 6, n_2 = 12, n_3 = 5, n_4 = 20$
60. A ball is drawn at random from box I and transferred to box II. If the probability of drawing a red ball from box I, after this transfer, is  $\frac{1}{3}$ , then the correct option(s) with the possible values of  $n_1$  and  $n_2$  is(are)
- (A)  $n_1 = 4, n_2 = 6$  (B)  $n_1 = 2, n_2 = 3$   
(C)  $n_1 = 10, n_2 = 20$  (D)  $n_1 = 3, n_2 = 6$



---

**(PAPER-2)**  
**JEE (ADVANCED) 2015**  
**ANSWERS**

**PART-I: PHYSICS**

- |     |             |     |             |     |             |     |                     |
|-----|-------------|-----|-------------|-----|-------------|-----|---------------------|
| 1.  | <b>2</b>    | 2.  | <b>7</b>    | 3.  | <b>4</b>    | 4.  | <b>6</b>            |
| 5.  | <b>3</b>    | 6.  | <b>2</b>    | 7.  | <b>2</b>    | 8.  | <b>1</b>            |
| 9.  | <b>A</b>    | 10. | <b>A, D</b> | 11. | <b>A, C</b> | 12. | <b>D</b>            |
| 13. | <b>A, B</b> | 14. | <b>B, C</b> | 15. | <b>D</b>    | 16. | <b>B or A, B, C</b> |
| 17. | <b>A, C</b> | 18. | <b>D</b>    | 19. | <b>A, D</b> | 20. | <b>A, C</b>         |

**PART-II: CHEMISTRY**

- |     |             |     |             |     |          |     |                |
|-----|-------------|-----|-------------|-----|----------|-----|----------------|
| 21. | <b>8</b>    | 22. | <b>4</b>    | 23. | <b>4</b> | 24. | <b>3</b>       |
| 25. | <b>5</b>    | 26. | <b>6</b>    | 27. | <b>3</b> | 28. | <b>9</b>       |
| 29. | <b>C</b>    | 30. | <b>A</b>    | 31. | <b>B</b> | 32. | <b>A</b>       |
| 33. | <b>B, C</b> | 34. | <b>C, D</b> | 35. | <b>B</b> | 36. | <b>B, C, D</b> |
| 37. | <b>A</b>    | 38. | <b>B</b>    | 39. | <b>C</b> | 40. | <b>D</b>       |

**PART-III: MATHEMATICS**

- |     |                |     |             |     |                |     |             |
|-----|----------------|-----|-------------|-----|----------------|-----|-------------|
| 41. | <b>9</b>       | 42. | <b>4</b>    | 43. | <b>9</b>       | 44. | <b>8</b>    |
| 45. | <b>4</b>       | 46. | <b>2</b>    | 47. | <b>9</b>       | 48. | <b>7</b>    |
| 49. | <b>D</b>       | 50. | <b>A, D</b> | 51. | <b>B, C, D</b> | 52. | <b>A, B</b> |
| 53. | <b>A, B, D</b> | 54. | <b>A, C</b> | 55. | <b>B, C</b>    | 56. | <b>A, B</b> |
| 57. | <b>A, B, C</b> | 58. | <b>C, D</b> | 59. | <b>A, B</b>    | 60. | <b>C, D</b> |

## SOLUTIONS

### PART-I: PHYSICS

1.  $mvr = \frac{nh}{2\pi} = \frac{3h}{2\pi}$

de-Broglie Wavelength  $\lambda = \frac{h}{mv} = \frac{2\pi r}{3} = \frac{2\pi a_0(3)^2}{3 z_{Li}} = 2\pi a_0$

2. For m closer to M

$$\frac{GMm}{9\ell^2} - \frac{Gm^2}{\ell^2} = ma \quad \dots(i)$$

and for the other m :

$$\frac{Gm^2}{\ell^2} + \frac{GMm}{16\ell^2} = ma \quad \dots(ii)$$

From both the equations,

$$k = 7$$

3.  $E(t) = A^2 e^{-\alpha t}$

$$\Rightarrow dE = -\alpha A^2 e^{-\alpha t} dt + 2AdAe^{-\alpha t}$$

Putting the values for maximum error,

$$\Rightarrow \frac{dE}{E} = \frac{4}{100} \Rightarrow \% \text{ error} = 4$$

4.  $I = \int \frac{2}{3} \rho 4\pi r^2 r^2 dr$

$$I_A \propto \int (r)(r^2)(r^2) dr$$

$$I_B \propto \int (r^5)(r^2)(r^2) dr$$

$$\therefore \frac{I_B}{I_A} = \frac{6}{10}$$

5. First and fourth wave interfere destructively. So from the interference of 2<sup>nd</sup> and 3<sup>rd</sup> wave only,

$$\Rightarrow I_{\text{net}} = I_0 + I_0 + 2\sqrt{I_0}\sqrt{I_0} \cos\left(\frac{2\pi}{3} - \frac{\pi}{3}\right) = 3I_0$$

$$\Rightarrow n = 3$$

6.  $\lambda_P = \frac{1}{\tau}; \lambda_Q = \frac{1}{2\tau}$

$$\frac{R_P}{R_Q} = \frac{(A_0 \lambda_P) e^{-\lambda_P t}}{A_0 \lambda_Q e^{-\lambda_Q t}}$$

$$\text{At } t = 2\tau; \frac{R_P}{R_Q} = \frac{2}{e}$$

7. Snell's Law on 1<sup>st</sup> surface :  $\frac{\sqrt{3}}{2} = n \sin r_1$

$$\sin r_1 = \frac{\sqrt{3}}{2n} \quad \dots(i)$$

$$\Rightarrow \cos r_1 = \sqrt{1 - \frac{3}{4n^2}} = \frac{\sqrt{4n^2 - 3}}{2n}$$

$$r_1 + r_2 = 60^\circ \quad \dots(ii)$$

Snell's Law on 2<sup>nd</sup> surface :

$$n \sin r_2 = \sin \theta$$

Using equation (i) and (ii)

$$n \sin (60^\circ - r_1) = \sin \theta$$

$$n \left[ \frac{\sqrt{3}}{2} \cos r_1 - \frac{1}{2} \sin r_1 \right] = \sin \theta$$

$$\frac{d}{dn} \left[ \frac{\sqrt{3}}{4} (\sqrt{4n^2 - 3} - 1) \right] = \cos \theta \frac{d\theta}{dn}$$


$$\text{for } \theta = 60^\circ \text{ and } n = \sqrt{3}$$

$$\Rightarrow \frac{d\theta}{dn} = 2$$

8. Equivalent circuit :

$$R_{eq} = \frac{13}{2} \Omega$$

So, current supplied by cell = 1 A



9. Q value of reaction =  $(140 + 94) \times 8.5 - 236 \times 7.5 = 219 \text{ Mev}$   
 So, total kinetic energy of Xe and Sr =  $219 - 2 - 2 = 215 \text{ Mev}$   
 So, by conservation of momentum, energy, mass and charge, only option (A) is correct

10. From the given conditions,  $\rho_1 < \sigma_1 < \sigma_2 < \rho_2$

From equilibrium,  $\sigma_1 + \sigma_2 = \rho_1 + \rho_2$

$$V_p = \frac{2}{9} \left( \frac{\rho_1 - \sigma_2}{\eta_2} \right) g \text{ and } V_Q = \frac{2}{9} \left( \frac{\rho_2 - \sigma_1}{\eta_1} \right) g$$

$$\text{So, } \frac{|\vec{V}_p|}{|\vec{V}_Q|} = \frac{\eta_1}{\eta_2} \text{ and } \vec{V}_p \cdot \vec{V}_Q < 0$$

11.  $BI/c \equiv VI \Rightarrow \mu_0 I^2 c \equiv VI \Rightarrow \mu_0 I c = V$

$$\Rightarrow \mu_0^2 I^2 c^2 = V^2$$


$$\Rightarrow \mu_0 I^2 = \epsilon_0 V^2 \Rightarrow \epsilon_0 c V = I$$

12.  $\vec{E} = \frac{\rho}{3\epsilon_0} \overrightarrow{C_1 C_2}$


$C_1 \Rightarrow$  centre of sphere and  $C_2 \Rightarrow$  centre of cavity.

13.  $Y = \frac{\text{stress}}{\text{strain}}$   
 $\Rightarrow \frac{1}{Y} = \frac{\text{strain}}{\text{stress}} \Rightarrow \frac{1}{Y_P} > \frac{1}{Y_Q} \Rightarrow Y_P < Y_Q$

14.  $P(r) = K \left( 1 - \frac{r^2}{R^2} \right)$




15.  $C_{10} = \frac{4\epsilon_0 \frac{S}{2}}{d/2} = \frac{4\epsilon_0 S}{d}$   
 $C_{20} = \frac{2\epsilon_0 S}{d}, C_{30} = \frac{\epsilon_0 S}{d}$   
 $\frac{1}{C'_{10}} = \frac{1}{C_{10}} + \frac{1}{C_{30}} = \frac{d}{2\epsilon_0 S} \left[ 1 + \frac{1}{2} \right]$   
 $\Rightarrow C'_{10} = \frac{4\epsilon_0 S}{3d}$   
 $C_2 = C_{30} + C'_{10} = \frac{7\epsilon_0 S}{3d}$   
 $\frac{C_2}{C_1} = \frac{7}{3}$



16.  $P$  (pressure of gas) =  $P_1 + \frac{kx}{A}$   
 $W = \int P dV = P_1(V_2 - V_1) + \frac{kx^2}{2} = P_1(V_2 - V_1) + \frac{(P_2 - P_1)(V_2 - V_1)}{2}$   
 $\Delta U = nC_V \Delta T = \frac{3}{2}(P_2 V_2 - P_1 V_1)$   
 $Q = W + \Delta U$   
 Case I:  $\Delta U = 3P_1 V_1, W = \frac{5P_1 V_1}{4}, Q = \frac{17P_1 V_1}{4}, U_{\text{spring}} = \frac{P_1 V_1}{4}$   
 Case II:  $\Delta U = \frac{9P_1 V_1}{2}, W = \frac{7P_1 V_1}{3}, Q = \frac{41P_1 V_1}{6}, U_{\text{spring}} = \frac{P_1 V_1}{3}$

**Note:** A and C will be true after assuming pressure to the right of piston has constant value  $P_1$ .

17.  $\theta \geq c$   
 $\Rightarrow 90^\circ - r \geq c$   
 $\Rightarrow \sin(90^\circ - r) \geq \sin c$   
 $\Rightarrow \cos r \geq \sin c$   
 using  $\frac{\sin i}{\sin r} = \frac{n_1}{n_m}$  and  $\sin c = \frac{n_2}{n_1}$   
 we get,  $\sin^2 i_m = \frac{n_1^2 - n_2^2}{n_m^2}$



Putting values, we get, correct options as A & C



18. For total internal reflection to take place in both structures, the numerical aperture should be the least one for the combined structure & hence, correct option is D.



19.  $I_1 = I_2$   
 $\Rightarrow n_1 A_1 v_1 = n_2 A_2 v_2$   
 $\Rightarrow d_1 w_1 v_1 = d_2 w_2 v_2$   
 Now, potential difference developed across MK  
 $V = Bvw$   
 $\Rightarrow \frac{V_1}{V_2} = \frac{v_1 w_1}{v_2 w_2} = \frac{d_2}{d_1}$   
 & hence correct choice is A & D

20. As  $I_1 = I_2$   
 $n_1 w_1 d_1 v_1 = n_2 w_2 d_2 v_2$   
 Now,  $\frac{V_2}{V_1} = \frac{B_2 v_2 w_2}{B_1 v_1 w_1} = \left( \frac{B_2 w_2}{B_1 w_1} \right) \left( \frac{n_1 w_1 d_1}{n_2 w_2 d_2} \right) = \frac{B_2 n_1}{B_1 n_2}$   
 $\therefore$  Correct options are A & C


## PART-II: CHEMISTRY

21.  $[\text{Fe}(\text{C}_2\text{O}_4)(\text{H}_2\text{O})]^{2-} + \text{MnO}_4^{2-} + 8\text{H}^+ \longrightarrow \text{Mn}^{2+} + \text{Fe}^{3+} + 4\text{CO}_2 + 6\text{H}_2\text{O}$   
 So the ratio of rate of change of  $[\text{H}^+]$  to that of rate of change of  $[\text{MnO}_4^-]$  is 8.





24.




The number of Fe – C bonds is 3.

25.  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$   $\longrightarrow$  will show cis – trans isomerism  
 $[\text{CrCl}_2(\text{C}_2\text{O}_4)_2]^{3-}$   $\longrightarrow$  will show cis – trans isomerism  
 $[\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2]^+$   $\longrightarrow$  will show cis – trans isomerism  
 $[\text{Fe}(\text{CN})_4(\text{NH}_3)_2]^-$   $\longrightarrow$  will show cis – trans isomerism  
 $[\text{Co}(\text{en})_2(\text{NH}_3)\text{Cl}]^{2+}$   $\longrightarrow$  will show cis – trans isomerism  
 $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]^{2+}$   $\longrightarrow$  will **not** show cis – trans isomerism (Although it will show geometrical isomerism)

26.  $\text{B}_2\text{H}_6 + 6\text{MeOH} \longrightarrow 2\text{B}(\text{OMe})_3 + 6\text{H}_2$   
 1 mole of  $\text{B}_2\text{H}_6$  reacts with 6 mole of  $\text{MeOH}$  to give 2 moles of  $\text{B}(\text{OMe})_3$ .  
 3 mole of  $\text{B}_2\text{H}_6$  will react with 18 mole of  $\text{MeOH}$  to give 6 moles of  $\text{B}(\text{OMe})_3$

27.  $\text{HX} \rightleftharpoons \text{H}^+ + \text{X}^-$

$$K_a = \frac{[\text{H}^+][\text{X}^-]}{[\text{HX}]}$$



$$K_a = \frac{[\text{H}^+][\text{Y}^-]}{[\text{HY}]}$$

$$\Lambda_m \text{ for HX} = \Lambda_{m_1}$$

$$\Lambda_m \text{ for HY} = \Lambda_{m_2}$$

$$\Lambda_{m_1} = \frac{1}{10} \Lambda_{m_2}$$

$$K_a = C\alpha^2$$

$$K_{a_1} = C_1 \times \left( \frac{\Lambda_{m_1}}{\Lambda_{m_1}^0} \right)^2$$




$$K_{a_2} = C_2 \times \left( \frac{\Lambda_{m_2}}{\Lambda_{m_2}^0} \right)^2$$


$$\frac{K_{a_1}}{K_{a_2}} = \frac{C_1}{C_2} \times \left( \frac{\Lambda_{m_1}}{\Lambda_{m_2}} \right)^2 = \frac{0.01}{0.1} \times \left( \frac{1}{10} \right)^2 = 0.001$$

$$pK_{a_1} - pK_{a_2} = 3$$


28. In conversion of  ${}^{238}_{92}\text{U}$  to  ${}^{206}_{82}\text{Pb}$ ,  $8\alpha$  - particles and  $6\beta$  particles are ejected.  
 The number of gaseous moles initially = 1 mol  
 The number of gaseous moles finally =  $1 + 8\text{ mol}$ ; (1 mol from air and 8 mol of  ${}^4_2\text{He}$ )  
 So the ratio =  $9/1 = 9$

29. At large inter-ionic distances (because  $a \rightarrow 0$ ) the P.E. would remain constant.  
 However, when  $r \rightarrow 0$ ; repulsion would suddenly increase.







34.  $\text{Cu}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Bi}^{3+}$  give ppt with  $\text{H}_2\text{S}$  in presence of dilute  $\text{HCl}$ .



36. \* Adsorption of  $\text{O}_2$  on metal surface is exothermic.  
 \* During electron transfer from metal to  $\text{O}_2$  electron occupies  $\pi^*_{2p}$  orbital of  $\text{O}_2$ .  
 \* Due to electron transfer to  $\text{O}_2$  the bond order of  $\text{O}_2$  decreases hence bond length increases.






$$\text{pH} = 5 - \log 2 + \log \frac{1/2}{1/2}$$

$$\text{pH} = 4.7$$

39.  $\text{C}_8\text{H}_6 \longrightarrow = \text{double bond equivalent} = 8 + 1 - \frac{6}{2} = 6$



JEE

## PART-III: MATHEMATICS

41.  $\vec{s} = 4\vec{p} + 3\vec{q} + 5\vec{r}$   
 $\vec{s} = x(-\vec{p} + \vec{q} + \vec{r}) + y(\vec{p} - \vec{q} + \vec{r}) + z(-\vec{p} - \vec{q} + \vec{r})$   
 $\vec{s} = (-x + y - z)\vec{p} + (x - y - z)\vec{q} + (x + y + z)\vec{r}$   
 $\Rightarrow -x + y - z = 4$   
 $\Rightarrow x - y - z = 3$   
 $\Rightarrow x + y + z = 5$

On solving we get  $x = 4, y = \frac{9}{2}, z = -\frac{7}{2}$

$\Rightarrow 2x + y + z = 9$

42. 
$$\frac{\sum_{k=1}^{12} \left| e^{\frac{k\pi}{7}} \right| \left| e^{\frac{\pi}{7}} - 1 \right|}{\sum_{k=1}^3 \left| e^{i(4k-2)} \right| \left| e^{\frac{\pi}{7}} - 1 \right|} = \frac{12}{3} = 4$$

43. Let seventh term be 'a' and common difference be 'd'

Given  $\frac{S_7}{S_{11}} = \frac{6}{11} \Rightarrow a = 15d$

Hence,  $130 < 15d < 140$

$\Rightarrow d = 9$

44.  $x^9$  can be formed in 8 ways  
 i.e.  $x^9, x^{1+8}, x^{2+7}, x^{3+6}, x^{4+5}, x^{1+2+6}, x^{1+3+5}, x^{2+3+4}$  and coefficient in each case is 1  
 $\Rightarrow$  Coefficient of  $x^9 = 1 + 1 + 1 + \dots + 1 = 8$   
8 times

45. The equation of  $P_1$  is  $y^2 - 8x = 0$  and  $P_2$  is  $y^2 + 16x = 0$

Tangent to  $y^2 - 8x = 0$  passes through  $(-4, 0)$

$\Rightarrow 0 = m_1(-4) + \frac{2}{m_1} \Rightarrow \frac{1}{m_1^2} = 2$

Also tangent to  $y^2 + 16x = 0$  passes through  $(2, 0)$

$\Rightarrow 0 = m_2 \times 2 - \frac{4}{m_2} \Rightarrow m_2^2 = 2$

$\Rightarrow \frac{1}{m_1^2} + m_2^2 = 4$

46. 
$$\lim_{\alpha \rightarrow 0} \frac{e^{\cos(\alpha^n)} - e}{\alpha^m} = -\frac{e}{2}$$

$$\lim_{\alpha \rightarrow 0} \frac{e \left( e^{(\cos(\alpha^n) - 1)} - 1 \right) (\cos \alpha^n - 1)}{(\cos(\alpha^n) - 1) \alpha^m \alpha^{2n}} \alpha^{2n} = -\frac{e}{2} \text{ if and only if } 2n - m = 0$$

$$47. \quad \alpha = \int_0^1 e^{(9x+3\tan^{-1}x)} \left( \frac{12+9x^2}{1+x^2} \right) dx$$

$$\text{Put } 9x + 3 \tan^{-1} x = t$$

$$\Rightarrow \left( 9 + \frac{3}{1+x^2} \right) dx = dt$$

$$\Rightarrow \alpha = \int_0^{9+\frac{3\pi}{4}} e^t dt = e^{9+\frac{3\pi}{4}} - 1$$

$$\Rightarrow \left( \log_e |1+\alpha| - \frac{3\pi}{4} \right) = 9$$

$$48. \quad G(1) = \int_{-1}^1 t |f(f(t))| dt = 0$$

$$f(-x) = -f(x)$$

$$\text{Given } f(1) = \frac{1}{2}$$

$$\lim_{x \rightarrow 1} \frac{F(x)}{G(x)} = \lim_{x \rightarrow 1} \frac{\frac{F(x)-F(1)}{x-1}}{\frac{G(x)-G(1)}{x-1}} = \frac{f(1)}{|f(f(1))|} = \frac{1}{14}$$

$$\Rightarrow \frac{1/2}{|f(1/2)|} = \frac{1}{14}$$

$$\Rightarrow f\left(\frac{1}{2}\right) = 7.$$

$$49. \quad \frac{192}{3} \int_{1/2}^x t^3 dt \leq f(x) \leq \frac{192}{2} \int_{1/2}^x t^3 dt$$

$$16x^4 - 1 \leq f(x) \leq 24x^4 - \frac{3}{2}$$

$$\int_{1/2}^1 (16x^4 - 1) dx \leq \int_{1/2}^1 f(x) dx \leq \int_{1/2}^1 \left( 24x^4 - \frac{3}{2} \right) dx$$

$$1 < \frac{26}{10} \leq \int_{1/2}^1 f(x) dx \leq \frac{39}{10} < 12$$

$$50. \quad \text{Here, } 0 < (x_1 - x_2)^2 < 1$$

$$\Rightarrow 0 < (x_1 + x_2)^2 - 4x_1x_2 < 1$$

$$\Rightarrow 0 < \frac{1}{\alpha^2} - 4 < 1$$

$$\Rightarrow \alpha \in \left( -\frac{1}{2}, -\frac{1}{\sqrt{5}} \right) \cup \left( \frac{1}{\sqrt{5}}, \frac{1}{2} \right)$$

51.  $\frac{\pi}{2} < \alpha < \pi, \pi < \beta < \frac{3\pi}{2} \Rightarrow \frac{3\pi}{2} < \alpha + \beta < \frac{5\pi}{2}$   
 $\Rightarrow \sin \beta < 0; \cos \alpha < 0$   
 $\Rightarrow \cos(\alpha + \beta) > 0.$

52. For the given line, point of contact for  $E_1 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  $\left(\frac{a^2}{3}, \frac{b^2}{3}\right)$

and for  $E_2 : \frac{x^2}{B^2} + \frac{y^2}{A^2} = 1$  is  $\left(\frac{B^2}{3}, \frac{A^2}{3}\right)$

Point of contact of  $x + y = 3$  and circle is  $(1, 2)$

Also, general point on  $x + y = 3$  can be taken as  $\left(1 \mp \frac{r}{\sqrt{2}}, 2 \pm \frac{r}{\sqrt{2}}\right)$  where,  $r = \frac{2\sqrt{2}}{3}$

So, required points are  $\left(\frac{1}{3}, \frac{8}{3}\right)$  and  $\left(\frac{5}{3}, \frac{4}{3}\right)$

Comparing with points of contact of ellipse,

$$a^2 = 5, B^2 = 8$$

$$b^2 = 4, A^2 = 1$$

$$\therefore e_1 e_2 = \frac{\sqrt{7}}{2\sqrt{10}} \text{ and } e_1^2 + e_2^2 = \frac{43}{40}$$

53. Tangent at P,  $xx_1 - yy_1 = 1$  intersects x axis at  $M\left(\frac{1}{x_1}, 0\right)$

$$\text{Slope of normal} = -\frac{y_1}{x_1} = \frac{y_1 - 0}{x_1 - x_2}$$

$$\Rightarrow x_2 = 2x_1 \Rightarrow N \equiv (2x_1, 0)$$

$$\text{For centroid } \ell = \frac{3x_1 + \frac{1}{x_1}}{3}, m = \frac{y_1}{3}$$

$$\frac{d\ell}{dx_1} = 1 - \frac{1}{3x_1^2}$$

$$\frac{dm}{dy_1} = \frac{1}{3}, \frac{dm}{dx_1} = \frac{1}{3} \frac{dy_1}{dx_1} = \frac{x_1}{3\sqrt{x_1^2 - 1}}$$

54. Let  $\int_0^\pi e^t (\sin^6 at + \cos^4 at) dt = A$

$$I = \int_\pi^{2\pi} e^t (\sin^6 at + \cos^4 at) dt$$

$$\text{Put } t = \pi + x$$

$$dt = dx$$

for  $a = 2$  as well as  $a = 4$

$$I = e^\pi \int_0^\pi e^x (\sin^6 ax + \cos^4 ax) dx$$

$$I = e^\pi A$$

$$\text{Similarly } \int_{2\pi}^{3\pi} e^t (\sin^6 at + \cos^4 at) dt = e^{2\pi} A$$

$$\text{So, } L = \frac{A + e^\pi A + e^{2\pi} A + e^{3\pi} A}{A} = \frac{e^{4\pi} - 1}{e^\pi - 1}$$

For both  $a = 2, 4$

55. Let  $H(x) = f(x) - 3g(x)$   
 $H(-1) = H(0) = H(2) = 3$ .  
 Applying Rolle's Theorem in the interval  $[-1, 0]$   
 $H'(x) = f'(x) - 3g'(x) = 0$  for atleast one  $c \in (-1, 0)$ .  
 As  $H''(x)$  never vanishes in the interval  
 $\Rightarrow$  Exactly one  $c \in (-1, 0)$  for which  $H'(x) = 0$   
 Similarly, apply Rolle's Theorem in the interval  $[0, 2]$ .  
 $\Rightarrow H'(x) = 0$  has exactly one solution in  $(0, 2)$

56.  $f(x) = (7\tan^6 x - 3\tan^2 x)(\tan^2 x + 1)$   
 $\int_0^{\pi/4} f(x) dx = \int_0^{\pi/4} (7\tan^6 x - 3\tan^2 x)\sec^2 x dx$   
 $\Rightarrow \int_0^{\pi/4} f(x) dx = 0$   
 $\int_0^{\pi/4} xf(x) dx = \left[ x \int f(x) dx \right]_0^{\pi/4} - \int_0^{\pi/4} \left[ \int f(x) dx \right] dx$   
 $\int_0^{\pi/4} xf(x) dx = \frac{1}{12}$

57. (A)  $f'(x) = F(x) + xF'(x)$   
 $f'(1) = F(1) + F'(1)$   
 $f'(1) = F'(1) < 0$   
 $f'(1) < 0$   
 (B)  $f(2) = 2F(2)$   
 $F(x)$  is decreasing and  $F(1) = 0$   
 Hence  $F(2) < 0$   
 $\Rightarrow f(2) < 0$   
 (C)  $f'(x) = F(x) + xF'(x)$   
 $F(x) < 0 \forall x \in (1, 3)$   
 $F'(x) < 0 \forall x \in (1, 3)$   
 Hence  $f'(x) < 0 \forall x \in (1, 3)$

58.  $\int_1^3 f(x) dx = \int_1^3 xF(x) dx$   
 $= \left[ \frac{x^2}{2} F(x) \right]_1^3 - \frac{1}{2} \int_1^3 x^2 F'(x) dx$   
 $= \frac{9}{2} F(3) - \frac{1}{2} F(1) + 6 = -12$   
 $40 = \left[ x^3 F'(x) \right]_1^3 - 3 \int_1^3 x^2 F'(x) dx$   
 $40 = 27F'(3) - F'(1) + 36 \quad \dots (i)$   
 $f'(x) = F(x) + xF'(x)$   
 $f'(3) = F(3) + 3F'(3)$   
 $f'(1) = F(1) + F'(1)$   
 $9f'(3) - f'(1) + 32 = 0$

59.  $P(\text{Red Ball}) = P(I) \cdot P(R | I) + P(II) \cdot P(R | II)$   
 $P(II | R) = \frac{1}{3} = \frac{P(II) \cdot P(R | II)}{P(I) \cdot P(R | I) + P(II) \cdot P(R | II)}$

$$\frac{1}{3} = \frac{\frac{n_3}{n_3 + n_4}}{\frac{n_1}{n_1 + n_2} + \frac{n_3}{n_3 + n_4}}$$

Of the given options, A and B satisfy above condition

60.  $P(\text{Red after Transfer}) = P(\text{Red Transfer}) \cdot P(\text{Red Transfer in II Case})$   
 $+ P(\text{Black Transfer}) \cdot P(\text{Red Transfer in II Case})$

$$P(R) = \frac{n_1}{n_1 + n_2} \cdot \frac{(n_1 - 1)}{(n_1 + n_2 - 1)} + \frac{n_2}{n_1 + n_2} \cdot \frac{n_1}{n_1 + n_2 - 1} = \frac{1}{3}$$

Of the given options, option C and D satisfy above condition.

