

PHYSICS

TEST PAPER WITH SOLUTION

SECTION-A

1. Match List I with List II :

List-I (Physical Quantity)		List-II (Dimensional Formula)	
A	Pressure gradient	I	$[M^0L^2T^{-2}]$
B	Energy density	II	$[M^1L^{-1}T^{-2}]$
C	Electric Field	III	$[M^1L^{-2}T^{-2}]$
D	Latent heat	IV	$[M^1L^1T^{-3}A^{-1}]$

Choose the **correct** answer from the options given below:

- (1) A-III, B-II, C-I, D-IV
 (2) A-II, B-III, C-IV, D-I
 (3) A-III, B-II, C-IV, D-I
 (4) A-II, B-III, C-I, D-IV

Official Ans. by NTA (3)

Allen Ans. (3)

Sol. Pressure gradient = $\frac{dp}{dx} = \frac{[ML^{-1}T^{-2}]}{[L]}$

= $[M^1L^{-2}T^{-2}]$

Energy density = $\frac{\text{energy}}{\text{volume}} = \frac{[ML^2T^{-2}]}{[L^3]}$

= $[M^1L^{-1}T^{-2}]$

Electric field = $\frac{\text{Force}}{\text{charge}} = \frac{[MLT^{-2}]}{[A.T]}$

= $[M^1L^1T^{-3}A^{-1}]$

Latent heat = $\frac{\text{heat}}{\text{mass}} = \frac{[ML^2T^{-2}]}{[M]}$

= $[M^0L^2T^{-2}]$

2. In a cuboid of dimension $2L \times 2L \times L$, a charge q is placed at the centre of the surface 'S' having area of $4L^2$. The flux through the opposite surface to 'S' is given by


(1) $\frac{q}{12\epsilon_0}$ (2) $\frac{q}{3\epsilon_0}$

(3) $\frac{q}{2\epsilon_0}$ (4) $\frac{q}{6\epsilon_0}$

Official Ans. by NTA (4)

Allen Ans. (4)

Sol. $\phi = \frac{Q/\epsilon_0}{6}$



Flux passing through shaded face = $\frac{q}{6\epsilon_0}$

3. Ratio of thermal energy released in two resistor R and 3R connected in parallel in an electric circuit is :

- (1) 3 : 1 (2) 1 : 1
 (3) 1 : 3 (4) 1 : 27




Official Ans. by NTA (1)

Allen Ans. (1)

Sol. $H = \frac{V^2}{R} \times t$

$\frac{H_1}{H_2} = \frac{\frac{V^2t}{R}}{\frac{V^2t}{3R}} = 3 : 1$


4. A single current carrying loop of wire carrying current I flowing in anticlockwise direction seen from +ve z direction and lying in xy plane in shown in figure. The plot of \hat{j} component of magnetic field (B_y) at a distance 'a' (less than radius of the coil) and on yz plane vs z coordinate look like



Official Ans. by NTA (3)

Allen Ans. (3)


Sol.



$B_y = 0$ in plane of coil

B_y is opposite of each other in $-z$ and $+z$ positions.

5. The magnitude of magnetic induction at mid-point O due to current arrangement as shown in Fig will be :



- (1) $\frac{\mu_0 I}{2\pi a}$ (2) 0
 (3) $\frac{\mu_0 I}{4\pi a}$ (4) $\frac{\mu_0 I}{\pi a}$


Official Ans. by NTA (4)

Allen Ans. (4)

- Sol. Magnetic field due to current in BC and ET are outward at point 'O'

$$B_0 = \frac{\mu_0 i}{4\pi r} + \frac{\mu_0 i}{4\pi r} = \frac{\mu_0 i}{2\pi r} = \frac{\mu_0 i}{\pi a}$$

6. Find the mutual inductance in the arrangement, when a small circular loop of wire of radius 'R' is placed inside a large square loop of wire of side L ($L \gg R$). The loops are coplanar and their centres coincide :



- (1) $M = \frac{\sqrt{2}\mu_0 R^2}{L}$ (2) $M = \frac{2\sqrt{2}\mu_0 R}{L^2}$
 (3) $M = \frac{2\sqrt{2}\mu_0 R^2}{L}$ (4) $M = \frac{\sqrt{2}\mu_0 R}{L^2}$

Official Ans. by NTA (3)

Allen Ans. (3)

Sol. $\phi = Mi$

$$\phi = (\mathbf{BA})$$

$$\phi = \pi R^2 \left(4 \frac{\mu_0}{4\pi} \frac{i}{\left(\frac{L}{2}\right)} \sqrt{2} \right)$$

$$\Rightarrow M = \frac{2\sqrt{2}\mu_0 R^2}{L}$$

7. Which of the following are true?

- A. Speed of light in vacuum is dependent on the direction of propagation.
 B. Speed of light in a medium is independent of the wavelength of light.
 C. The speed of light is independent of the motion of the source.
 D. The speed of light in a medium is independent of intensity.

Choose the correct answer from the option given below :


- (1) A and C only (2) B and D only
 (3) B and C only (4) C and D only

Official Ans. by NTA (4)

Allen Ans. (4)

- Sol. Speed of light does not depend on the motion of source as well as intensity.

8. In a Young's double slit experiment, two slits are illuminated with a light of wavelength 800 nm. The line joining A_1P is perpendicular to A_1A_2 as shown in the figure. If the first minimum is detected at P, the value of slits separation 'a' will be :



The distance of screen from slits $D = 5$ cm

- (1) 0.4 mm (2) 0.5 mm
 (3) 0.2 mm (4) 0.1 mm

Official Ans. by NTA (3)

Allen Ans. (3)

26. A tennis ball is dropped on to the floor from a height of 9.8 m. It rebounds to a height 5.0 m. Ball comes in contact with the floor for 0.2s. The average acceleration during contact is _____ ms^{-2} .
[Given $g = 10 \text{ ms}^{-2}$]

Official Ans. by NTA (120)

Allen Ans. (120)

Sol.

$$v_i = \sqrt{2gh_i}$$

$$= \sqrt{2 \times 10 \times 9.8} \downarrow$$

$$= 14 \text{ m/s} \downarrow$$

$$v_f = \sqrt{2gh_f}$$

$$= \sqrt{2 \times 10 \times 5} \uparrow$$


$$= 10 \text{ m/s} \uparrow$$

$$|\bar{a}_{\text{avg}}| = \left| \frac{\Delta \bar{v}}{\Delta t} \right| = \frac{24}{0.2} = 120 \text{ m/s}^2$$

27. A point charge $q_1 = 4q_0$ is placed at origin. Another point charge $q_2 = -q_0$ is placed at $x = 12 \text{ cm}$. Charge of proton is q_0 . The proton is placed on x-axis so that the electrostatic force on the proton is zero. In this situation, the position of the proton from the origin is _____ cm.

Official Ans. by NTA (24)

Allen Ans. (24)



$$\frac{q_0}{x^2} = \frac{4q_0}{(x+12)^2}$$

$$x+12 = 2x$$

$$x = 12$$

Distance from origin = $x + 12 = 24 \text{ cm}$.

28. In a metre bridge experiment the balance point is obtained if the gaps are closed by 2Ω and 3Ω . A shunt of $X\Omega$ is added to 3Ω resistor to shift the balancing point by 22.5 cm. The value of X is _____

Official Ans. by NTA (2)

Allen Ans. (2)

Sol.

$$\frac{2}{\left(\frac{3x}{3+x}\right)} = \frac{40+22.5}{60-22.5} = \frac{62.5}{37.5} = \frac{5}{3}$$

$$\frac{6}{5} = \frac{3x}{3+x}$$

$$6+2x = 5x \Rightarrow x = 2$$

29. A certain elastic conducting material is stretched into a circular loop. It is placed with its plane perpendicular to a uniform magnetic field $B = 0.8 \text{ T}$. When released the radius of the loop starts shrinking at a constant rate of 2 cm^{-1} . The induced emf in the loop at an instant when the radius of the loop is 10 cm will be _____ mV.

Official Ans. by NTA (10)

Allen Ans. (10)

Sol.


$$\text{EMF} = \frac{d}{dt}(B\pi r^2)$$

$$= 2B\pi r \frac{dr}{dt} = 2 \times \pi \times 0.1 \times 0.8 \times 2 \times 10^{-2}$$

$$= 2\pi \times 1.6 = 10.06 \text{ [round off } 10.06 = 10]$$

30. As shown in figures, three identical polaroids P_1 , P_2 and P_3 are placed one after another. The pass axis of P_2 and P_3 are inclined at angle of 60° and 90° with respect to axis of P_1 . The source S has an intensity of $256 \frac{\text{W}}{\text{m}^2}$. The intensity of light at point

O is _____ $\frac{\text{W}}{\text{m}^2}$.



Official Ans. by NTA (24)

Allen Ans. (24)

- Sol.** By first polaroid P_1 intensity will be halved then P_2 and P_3 will make intensity $\cos^2(60^\circ)$ and $\cos^2(30^\circ)$ times respectively.

$$\text{Intensity out} = \frac{256}{2} \times \frac{1}{4} \times \left(\frac{\sqrt{3}}{2}\right)^2 = \frac{256 \times 3}{2 \times 4 \times 4} = 24$$