

CHEMISTRY
JEE-MAIN (February-Attempt) 25
February (Shift-1) Paper

SECTION - A

1. Ellingham diagram is a graphical representation of:
 (1) ΔG vs T (2) $(\Delta G - T\Delta S)$ vs T (3) ΔH vs T (4) ΔG vs P

Sol. (1)
 Ellingham diagram tells us about the spontaneity of a reaction with temperature.

2. Which of the following equation depicts the oxidizing nature of H_2O_2 ?
 (1) $Cl_2 + H_2O_2 \rightarrow 2HCl + O_2$ (2) $KIO_4 + H_2O_2 \rightarrow KIO_3 + H_2O + O_2$
 (3) $2I^- + H_2O_2 + 2H^+ \rightarrow I_2 + 2H_2O$ (4) $I_2 + H_2O_2 + 2OH^- \rightarrow 2I^- + 2H_2O + O_2$

Sol. (3)
 $2I^- + H_2O_2 + 2H^+ \rightarrow I_2 + 2H_2O$
 Oxygen reduces from -1 to -2
 So, its reduction will take place. Hence it will behave as oxidising agent or it shows oxidising nature.
 While in other option it change from (-1) to 0.

3. In Freundlich adsorption isotherm at moderate pressure, the extent of adsorption $\left(\frac{x}{m}\right)$ is directly proportional to P^x . The value of x is:

- (1) ∞ (2) 1 (3) zero (4) $\frac{1}{n}$

Sol. (4)

$$\frac{x}{m} = p^x$$

the formula is $\frac{x}{m} = p^{\frac{1}{n}}$


Hence $x = \frac{1}{n}$

The value of 'n' is any natural number.

4. According to molecular orbital theory, the species among the following that does not exist is:
 (1) He_2^- (2) He_2^+ (3) O_2^{2-} (4) Be_2

Sol. (4)
 B.O. of Be_2 is zero, So it does not exist.


5. Identify A in the given chemical reaction.



Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

Sol. (4)



Aromatization reaction or hydroforming reaction.


6. Given below are two statements:
Statement-I : CeO_2 can be used for oxidation of aldehydes and ketones.
Statement-II : Aqueous solution of EuSO_4 is a strong reducing agent.

- (1) Statement I is true, statement II is false
(2) Statement I is false, statement II is true
(3) Both Statement I and Statement II are false
(4) Both Statement I and Statement II are true

Sol. (4)


CeO_2 can be used as oxidising agent like seO_2 .
Similarly EuSO_4 used as a reducing agent.

7. The major product of the following chemical reaction is:



- (1) $(\text{CH}_3\text{CH}_2\text{CO})_2\text{O}$ (2) $\text{CH}_3\text{CH}_2\text{CHO}$ (3) $\text{CH}_3\text{CH}_2\text{CH}_3$ (4) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$

Sol. (2)



8. Complete combustion of 1.80 g of an oxygen containing compound ($\text{C}_x\text{H}_y\text{O}_z$) gave 2.64 g of CO_2 and 1.08 g of H_2O . The percentage of oxygen in the organic compound is:
(1) 63.53 (2) 53.33 (3) 51.63 (4) 50.33

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

Sol. (2)

$$n_{\text{CO}_2} = \frac{2.64}{44} = 0.06$$

$$n_c = 0.06$$

$$\text{weight of carbon} = 0.06 \times 12 = 0.72 \text{ gm}$$

$$n_{\text{H}_2\text{O}} = \frac{1.08}{18} = 0.06$$

$$n_H = 0.06 \times 2 = 0.12$$

$$\text{weight of H} = 0.12 \text{ gm}$$

$$\therefore \text{Weight of oxygen in } C_xH_yO_z$$

$$= 1.8 - (0.72 + 0.12)$$

$$= 0.96 \text{ gram}$$

$$\begin{aligned} \% \text{ weight of oxygen} &= \frac{0.96}{1.8} \times 100 \\ &= 53.3\% \end{aligned}$$

- 9.** The correct statement about B_2H_6 is:
- (1) All B-H-B angles are of 120° .
 - (2) Its fragment, BH_3 , behaves as a Lewis base.
 - (3) Terminal B-H bonds have less p-character when compared to bridging bonds.
 - (4) The two B-H-B bonds are not of same length.

Sol. (3)



Terminal bond angle is greater than that of bridge bond angle

Bond angle \propto s-character

$$\propto \frac{1}{p - \text{character}}$$

- 10.** In which of the following pairs, the outer most electronic configuration will be the same?
- (1) Fe^{2+} and Co^+
 - (2) Cr^+ and Mn^{2+}
 - (3) Ni^{2+} and Cu^+
 - (4) V^{2+} and Cr^+

Sol. (2)



- 11.** Which statement is correct?
- (1) Buna-S is a synthetic and linear thermosetting polymer
 - (2) Neoprene is addition copolymer used in plastic bucket manufacturing
 - (3) Synthesis of Buna-S needs nascent oxygen
 - (4) Buna-N is a natural polymer

Sol. (3)

Synthesis of Buna-S needs nascent oxygen.





Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

12. Given below are two statements:
Statement-I : An allotrope of oxygen is an important intermediate in the formation of reducing smog.
Statement-II : Gases such as oxides of nitrogen and sulphur present in troposphere contribute to the formation of photochemical smog.
In the light of the above statements, choose the correct answer from the options given below:
(1) Statement I and Statement II are true
(2) Statement I is true about Statement II is false
(3) Both Statement I and Statement II are false
(4) Statement I is false but Statement II is true

Sol. (3)
Reducing smog as is acts as reducing agent, the reducing character is due to presence of sulphur dioxide and carbon particles.

13. The plots of radial distribution functions for various orbitals of hydrogen atom against 'r' are given below:



The correct plot for 3s orbital is:

- (1) D (2) B (3) A (4) C

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

Sol. (1)

3s orbital

Number of radial nodes = $n - \ell - 1$

For 3s orbital $n = 3$ $\ell = 0$


Number of radial nodes = $3 - 0 - 1 = 2$

It is correctly represented in graph of option D

14. Which of the glycosidic linkage galactose and glucose is present in lactose?

- (1) C-1 of glucose and C-6 of galactose (2) C-1 of galactose and C-4 of glucose
 (3) C-1 of glucose and C-4 of galactose (4) C-1 of galactose and C-6 of glucose





Sol. (2)




β -D-Galactose

β -D-Glucose

15. Which one of the following reactions will not form acetaldehyde?






Sol. (1)



Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

16. Which of the following reaction/s will not give p-aminoazobenzene?




(1) B only

(2) A and B

(3) C only

(4) A only

Sol. (1)



17. The hybridization and magnetic nature of $[\text{Mn}(\text{CN})_6]^{4-}$ and $[\text{Fe}(\text{CN})_6]^{3-}$, respectively are:



(1) d^2sp^3 and paramagnetic

(2) sp^3d^2 and paramagnetic

(3) d^2sp^3 and diamagnetic

(4) sp^3d^2 and diamagnetic






Sol. (1)




Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

18. Identify A and B in the chemical reaction.




Sol. (4)



Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

19. Compound(s) which will liberate carbon dioxide with sodium bicarbonate solution is/are:



- (1) B and C only (2) B only (3) A and B only (4) C only

Sol. (1)

Compounds which are more acidic than H_2CO_3 , gives CO_2 gas on reaction with NaHCO_3 . Compound B i.e. Benzoic acid and compound C i.e. picric acid both are more acidic than H_2CO_3 .

20. The solubility of AgCN in a buffer solution of $\text{pH} = 3$ is x . The value of x is:

[Assume: No cyano complex is formed; $K_{\text{sp}}(\text{AgCN}) = 2.2 \times 10^{-16}$ and $K_a(\text{HCN}) = 6.2 \times 10^{-10}$]

(1) 0.625×10^{-6} (2) 1.6×10^{-6} (3) 2.2×10^{-16} (4) 1.9×10^{-5}

Sol. (4)

Let solubility is x



$$K_{\text{sp}} \times \frac{1}{K_a} = [\text{Ag}^+][\text{CN}^-] \times \frac{[\text{HCN}]}{[\text{H}^+][\text{CN}^-]}$$

$$2.2 \times 10^{-16} \times \frac{1}{6.2 \times 10^{-10}} = \frac{[\text{S}][\text{S}]}{10^{-3}}$$

$$S^2 = \frac{2.2}{6.2} \times 10^{-9}$$

$$S^2 = 3.55 \times 10^{-10}$$

$$S = \sqrt{3.55 \times 10^{-10}}$$

$$S = 1.88 \times 10^{-5} \Rightarrow 1.9 \times 10^{-5}$$


SECTION - B

1. The reaction of cyanamide, $\text{NH}_2\text{CN}_{(s)}$ with oxygen was run in a bomb calorimeter and ΔU was found to be $-742.24 \text{ kJ mol}^{-1}$. The magnitude of ΔH_{298} for the reaction



[Assume ideal gases and $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$]

Sol. 741 kJ/mol



$$\Delta n_g = (1 + 1) - \frac{3}{2} = \frac{1}{2}$$


Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

$$\begin{aligned}\Delta H &= \Delta U + \Delta ng RT \\ &= -742.24 + \frac{1}{2} \times \frac{8.314 \times 298}{1000} \\ &= -742.24 + 1.24 \\ &= 741 \text{ kJ/mol}\end{aligned}$$

2. In basic medium CrO_4^{2-} oxidizes $\text{S}_2\text{O}_3^{2-}$ to form SO_4^{2-} and itself changes into $\text{Cr}(\text{OH})_4^-$. The volume of 0.154 M CrO_4^{2-} required to react with 40 mL of 0.25 M $\text{S}_2\text{O}_3^{2-}$ is _____ mL. (Rounded-off to the nearest integer)

Sol. 173 mL




Applying mole-mole analysis

$$\frac{0.154 \times v}{8} = \frac{40 \times 0.25}{3}$$

$$V = 173 \text{ mL}$$

3. For the reaction, $a\text{A} + b\text{B} \rightarrow c\text{C} + d\text{D}$, the plot of $\log k$ vs $\frac{1}{T}$ is given below:



The temperature at which the rate constant of the reaction is 10^{-4}s^{-1} is _____ K. [Rounded off to the nearest integer]

[Given: The rate constant of the reaction is 10^{-5}s^{-1} at 500 K]

Sol. 526 K

$$\log_{10}K = \log_{10}A - \frac{E_a}{2.303RT}$$

$$\text{Slope} = \frac{E_a}{2.303R} = -10000$$

$$\log_{10} \frac{K_2}{K_1} = \frac{E_a}{2.303R} \times \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$\log_{10} \frac{10^{-4}}{10^{-5}} = 10000 \times \left[\frac{1}{500} - \frac{1}{T} \right]$$

$$1 = 10000 \times \left[\frac{1}{500} - \frac{1}{T} \right]$$

$$\frac{1}{10000} = \frac{1}{500} - \frac{1}{T}$$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

$$\frac{1}{T} = \frac{1}{500} - \frac{1}{10000}$$


$$\frac{1}{T} = \frac{20 - 1}{10000} = \frac{19}{10000}$$

$$T = \frac{10,000}{19} \Rightarrow 526 \text{ K}$$


4. 0.4g mixture of NaOH, Na₂CO₃ and some inert impurities was first titrated with $\frac{N}{10}$ HCl using phenolphthalein as an indicator, 17.5 mL of HCl was required at the end point. After this methyl orange was added and titrated. 1.5 mL of same HCl was required for the next end point. The weight percentage of Na₂CO₃ in the mixture is _____. (Rounded-off to the nearest integer)

Sol. 3%

1st end point reaction



$$nf = 1$$




$$nf = 1$$

$$\text{Eq of HCl used} = n_{\text{NaOH}} \times 1 + n_{\text{Na}_2\text{CO}_3} \times 1$$

$$17.5 \times \frac{1}{10} \times 10^{-3} = n_{\text{NaOH}} + n_{\text{Na}_2\text{CO}_3}$$

2nd end point



$$1.5 \times \frac{1}{10} \times 10^{-3} = n_{\text{NaHCO}_3} \times 1 = n_{\text{NaHCO}_3}$$

$$0.15 \text{ mmol} = n_{\text{Na}_2\text{CO}_3}$$

$$0.15 = n_{\text{Na}_2\text{CO}_3}$$




$$W_{\text{Na}_2\text{CO}_3} = \frac{0.15 \times 106 \times 10^{-3}}{0.5} \times 100 \times 10$$

$$= 3 \times 106 \times 10^{-2}$$

$$= 3 \times 1.06 = 3.18\%$$

5. The ionization enthalpy of Na⁺ formation from Na_(g) is 495.8 kJ mol⁻¹, while the electron gain enthalpy of Br is -325.0 kJ mol⁻¹. Given the lattice enthalpy of NaBr is -728.4 kJ mol⁻¹. The energy for the formation of NaBr ionic solid is (-)_____ × 10⁻¹ kJ mol⁻¹.

Sol. 5576 kJ




$$\Delta H = 495.8 - 325 - 728.4$$

$$-557.6 \text{ kJ} = -5576 \times 10^{-1} \text{ kJ}$$

Toll Free : 1800-212-1799


www.motion.ac.in | Email : info@motion.ac.in

6. Consider the following chemical reaction.



The number of sp^2 hybridized carbon atom(s) present in the product is _____.

Sol. 7



All carbon atoms in benzaldehyde are sp^2 hybridised

7. A car tyre is filled with nitrogen gas at 35 psi at 27°C. It will burst if pressure exceeds 40 psi. The temperature in °C at which the car tyre will burst is _____. (Rounded-off to the nearest integer)

Sol. 69.85°C ≈ 70°C

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$\frac{35}{300} = \frac{40}{T_2}$$

$$T_2 = \frac{40 \times 300}{35}$$

$$= 342.86 \text{ K}$$

$$= 69.85^\circ\text{C} \approx 70^\circ\text{C}$$

8. Among the following, the number of halide(s) which is/are inert to hydrolysis is _____.

(A) BF_3 (B) SiCl_4 (C) PCl_5 (D) SF_6


Sol. 1

Due to crowding SF_6 is not hydrolysed.

9. 1 molal aqueous solution of an electrolyte A_2B_3 is 60% ionised. The boiling point of the solution at 1 atm is _____ K. (Rounded-off to the nearest integer)

[Given K_b for $(\text{H}_2\text{O}) = 0.52 \text{ K kg mol}^{-1}$]

Sol. 375 K



$$\text{No. of ions} = 2 + 3 = 5$$

$$i = 1 + (n - 1) \alpha$$

$$= 1 + (5 - 1) \times 0.6$$

$$= 1 + 4 \times 0.6 = 1 + 2.4 = 3.4$$

$$\Delta T_b = K_b \times m \times i$$

$$= 0.52 \times 1 \times 3.4 = 1.768^\circ\text{C}$$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

$$\Delta T_b = (T_b)_{\text{solution}} - [(T_b)_{\text{H}_2\text{O}}]_{\text{solution}}$$

$$1.768 = (T_b)_{\text{solution}} - 100$$

$$(T_b)_{\text{solution}} = 101.768 \text{ } ^\circ\text{C}$$

$$= 375 \text{ K}$$

10. Using the provided information in the following paper chromatogram:




Fig: Paper chromatography for compounds A and B
the calculated R_f value of A _____ $\times 10^{-1}$.

Sol. 4

$$R_f = \frac{\text{Distance travelled by compound}}{\text{Distance travelled by solvent}}$$

On chromatogram distance travelled by compound is $\rightarrow 2 \text{ cm}$

Distance travelled by solvent = 5 cm

$$\text{So } R_f = \frac{2}{5} = 4 \times 10^{-1} = 0.4$$