

MATHEMATICS

1. Find number of common terms in the two given series

4, 9, 14, 19..... up to 25 terms and

3, 9, 15, 21up to 37 terms

Ans. (1)

Sol. 4, 9, 14, 19, 124 $\rightarrow d_1 = 5$

$$3, 9, 15, 21 \dots\dots\dots 219 \rightarrow d_2 = 6$$

1st common term = 9 and common difference of common terms = 30

Common terms are 9, 39, 69, 99

4 common terms

2. Let $8 = 3 + \frac{3+p}{4} + \frac{3+2p}{4^2} + \dots \infty$, then p is

Ans. (1)

$$\text{Sol. } 8 = 3 + \frac{3+p}{4} + \frac{3+2p}{4^2} + \dots \dots \dots \text{(i)}$$

multiply both sides by $\frac{1}{4}$, we get

$$2 = \frac{3}{4} + \frac{3+p}{4^2} + \dots \quad (\text{ii})$$

Equation (i) – equation (ii)

$$\Rightarrow 6 = 3 + \frac{p}{4} + \frac{p}{4^2} + \dots$$

$$\Rightarrow 3 = \frac{p}{4\left(1 - \frac{1}{4}\right)} \Rightarrow p = 9$$

3. For $\frac{x^2}{25} + \frac{y^2}{16} = 1$, find the length of chord whose mid point is $P\left(1, \frac{2}{5}\right)$

- $$(1) \frac{\sqrt{1681}}{5} \quad (2) \frac{\sqrt{1481}}{5} \quad (3) \frac{\sqrt{1781}}{5} \quad (4) \frac{\sqrt{1691}}{5}$$

Ans. (4)

Sol. By $T = S_1$

$$\Rightarrow \frac{x}{25} + \frac{y}{16} = \frac{1}{25} + \frac{4}{25} \cdot \frac{1}{16}$$

$$\Rightarrow \frac{x}{25} + \frac{y}{40} = \frac{4+1}{100}$$

$$\Rightarrow \frac{x}{25} + \frac{y}{40} = \frac{1}{20}$$

$$\Rightarrow 8x + 5y = 10$$

$$\Rightarrow \frac{x^2}{25} + \left(\frac{10-8x}{5} \right)^2 \frac{1}{16} = 1$$

$$\Rightarrow \frac{x^2}{25} + \frac{4}{25} \left(\frac{5-4x}{16} \right)^2 = 1$$

$$\Rightarrow x^2 + \frac{(5-4x)^2}{4} = 25$$

$$\Rightarrow 4x^2 + (5-4x)^2 = 100$$

$$\Rightarrow 20x^2 - 8x - 15 = 0$$

$$x_1 + x_2 = 2$$

$$x_1 x_2 = \frac{-15}{4}$$

$$\text{length of chord} = |x_1 - x_2| \sqrt{1 + m^2}$$

$$= \frac{\sqrt{1691}}{5}$$

4. If $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$, then find $f'(10)$.

Ans. (202)

$$\text{Sol. } f'(x) = 3x^2 + 2xf'(1) + f''(2)$$

$$f''(x) = 6x + 2f'(1)$$

$$f'''(3) = 6$$

$$f'(1) = -5$$

$$f''(2) = 2$$

$$\Rightarrow f(10) = 300 + 20(-5) + 2$$

$$= 202$$

5. Let $\int_0^1 \frac{dx}{\sqrt{x+3} + \sqrt{x+1}} = A + B\sqrt{2} + C\sqrt{3}$ then the value of $2A + 3B + C$ is

(1) 3

(2) 4

(3) 5

(4) 6

Ans. (1)

Sol. On rationalising

$$\begin{aligned}
 & \int_0^1 \frac{(\sqrt{x+3} - \sqrt{x+1})}{2} dx \\
 &= \frac{2}{3.2} \left\{ (x+3)^{3/2} - (x+1)^{3/2} \right\}_0^1 \\
 &= \frac{1}{3} \{8 - 3\sqrt{3} - (2\sqrt{2} - 1)\} \\
 &= \frac{1}{3} \{9 - 3\sqrt{3} - 2\sqrt{2}\} \\
 &= \left(3 - \sqrt{3} - \frac{2\sqrt{2}}{3} \right) : A = 3, B = -\frac{2}{3}, C = -1
 \end{aligned}$$

$$\therefore 2A + 3B + C = 6 - 2 - 1 = 3$$

Ans. (2)

- Sol.** z is equidistant from 1, i, & -i
only $z = 0$ is possible
 \therefore number of z equal to 1

7. If sum of coefficients in $(1 - 3x + 10x^2)^n$ and $(1 + x^2)^n$ is A and B respectively then
 (1) $A^3 = B$ (2) $A = B^3$ (3) $A = 2B$ (4) $A = B$

Ans. (2)

- $$\textbf{Sol.} \quad A = 8^n \quad B = 2^n$$

$$(B) \therefore A \equiv B^3$$

8. Let a_1, a_2, \dots, a_{10} are 10 observations such that $\sum_{i=1}^{10} a_i = 50$ and $\sum_{i \neq j} a_i \cdot a_j = 1100$, then their standard deviation will be

Ans. (1)

$$\begin{aligned}
 \text{Sol. } & (a_1 + a_2 + \dots + a_{10})^2 = 50^2 \\
 & \Rightarrow \sum a_i^2 + 2 \sum_{i \neq j} a_i a_j = 2500 \\
 & \Rightarrow \sum a_i^2 = 300 \\
 & \sigma^2 = \frac{\sum a_i^2}{10} - \left(\frac{\sum a_i}{10} \right)^2 \\
 & \Rightarrow \sigma^2 = 5 \Rightarrow S.D = \sqrt{5}
 \end{aligned}$$

9. If $f(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$ then

Statement-1 : $f(-x)$ is inverse of $f(x)$

Statement-2 : $f(x + y) = f(x)f(y)$

Ans. (1)

$$\text{Sol. } f(x)f(y) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos y & -\sin y & 0 \\ \sin y & \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} \cos(x+y) & -\sin(x+y) & 0 \\ \sin(x+y) & \cos(x-y) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= f(x + y)$$

$$\therefore f(x) f(-x) = f(0)$$

= I

- 10.** If $a = \lim_{x \rightarrow 0} \frac{\sqrt{1 + \sqrt{1 + x^4}} - \sqrt{2}}{x^4}$ and $b = \lim_{x \rightarrow 0} \frac{\sin^2 x}{\sqrt{2} - \sqrt{1 + \cos x}}$ find $a \cdot b^3$

(1) 16 (2) 32 (3) -16 (4) 48

Ans. (2)

$$\text{Sol. } a = \lim_{x \rightarrow 0} \frac{\sqrt{1+x^4} - 1}{x^4 \left[\sqrt{1+\sqrt{1+x^4}} + \sqrt{2} \right]}$$

$$= \lim_{x \rightarrow 0} \frac{x^4}{x^4 \left[\sqrt{1 + \sqrt{1 + x^4}} + \sqrt{2} \right] \left[\sqrt{1 + x^4} + 1 \right]}$$

$$= \frac{1}{2\sqrt{2} \times 2} = \frac{1}{4\sqrt{2}}$$

$$b = \lim_{x \rightarrow 0} \frac{\sin^2 x}{(1 - \cos x)} \left(\sqrt{2} + \sqrt{1 + \cos x} \right)$$

$$= 2 \times (\sqrt{2} + \sqrt{2}) = 4\sqrt{2}$$

$$\therefore ab^3 = (4\sqrt{2})^2 = 32$$

11. If the minimum distance of centre of the circle $x^2 + y^2 - 4x - 16y + 64 = 0$ from any point on the parabola $y^2 = 4x$ is d , find d^2

Ans. (20)

Sol. Normal to parabola is $y = mx - 2m - m^3$

$$\text{centre } (2, 8) \rightarrow 8 = 2m - 2m - m^3$$

$$\Rightarrow m = -2$$

$$\therefore p \text{ is } (m^2, -2m) = (4, 4)$$

$$\Rightarrow d^2 = 4 + 16 = 20$$

12. If $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$, $\vec{b} = 3(\hat{i} - \hat{j} + \hat{k})$, $\vec{a} \times \vec{c} = \vec{b}$ & $\vec{a} \cdot \vec{c} = 3$ find $\vec{a} \cdot (\vec{c} \times \vec{b} - \vec{b} \cdot \vec{c})$

(1) 24

(2) -24

(3) 18

(4) 15

Ans. (1)

Sol. $[\vec{a} \vec{c} \vec{b}] = (\vec{a} \times \vec{c}) \cdot \vec{b} = |\vec{b}|^2 = 27$

$$\therefore \text{we need} = 27 - 0 - 3 = 24$$

13. Consider the line $L : 4x + 5y = 20$. Let two other lines are L_1 and L_2 which trisect the line L and pass through origin, then tangent of angle between lines L_1 and L_2 is

(1) $\frac{20}{41}$

(2) $\frac{30}{41}$

(3) $\frac{40}{41}$

(4) $\frac{10}{41}$

Ans. (2)

Sol. Let line L intersect the lines L_1 and L_2 at P and Q

$$P\left(\frac{10}{3}, \frac{4}{3}\right), Q\left(\frac{5}{3}, \frac{8}{3}\right)$$

$$\therefore m_{OA} = \frac{2}{5}$$

$$m_{OQ} = \frac{8}{5}$$

$$\tan \theta = \left| \frac{\frac{8}{5} - \frac{2}{5}}{\frac{16}{25}} \right|$$

$$= \left(\frac{6}{5} \times \frac{25}{41} \right)$$

$$= \frac{30}{41}$$

14. If ${}^{n-1}C_r = (k^2 - 8) {}^nC_{r+1}$, then the range of 'k' is

(1) $k \in (2\sqrt{2}, 3]$ (2) $k \in (2\sqrt{2}, 3)$ (3) $k \in [2, 3)$ (4) $k \in (2\sqrt{2}, 8)$

Ans. (1)

Sol. ${}^{n-1}C_r = (k^2 - 8) \frac{n}{r+1} \cdot {}^nC_r$

$$\Rightarrow k^2 - 8 = \frac{r+1}{n}$$

here $r \in [0, n-1]$

$$\Rightarrow r+1 \in [1, n]$$

$$\Rightarrow k^2 - 8 \in \left[\frac{1}{n}, 1 \right]$$

$$\Rightarrow k^2 \in \left[8 + \frac{1}{n}, 9 \right]$$

$$\Rightarrow k \in (2\sqrt{2}, 3]$$

15. If $\alpha x + \beta y + 9\ln|2x + 3y - 8| = x + C$ is the solution of $(2x + 3y - 2)dx + (4x + 6y - 7)dy = 0$, then $\alpha + \beta + \gamma =$

(1) 18

(2) 19

(3) 20

(4) 21

Ans. (1)

Sol. Let $2x + 3y = t$

$$\Rightarrow 2 + 3 \frac{dy}{dx} = \frac{dt}{dx}$$

Now $(t-2) + (2t-7) \left(\frac{dt}{dx} - 2 \right) \times \frac{1}{3} = 0$

$$\Rightarrow -\frac{(3t-6)}{2t-7} = \frac{dt}{dx} - 2$$

$$\Rightarrow \frac{dt}{dx} = \frac{t-8}{2t-7}$$

$$\Rightarrow \int \frac{2t-7}{t-8} dt = \int dx$$

$$\Rightarrow \int 2 + \frac{9}{t-8} dt = \int dx$$

$$\Rightarrow 2t + 9\ln|t-8| = x + C$$

$$\Rightarrow 2(2x + 3y) + 9\ln|2x + 3y - 8| = x + C$$

$$\alpha = 4, \beta = 6, \gamma = 8$$

Ans. (3)

Sol. '4' is not image of any element \Rightarrow into

$$f(10) = 5 = f(15) \Rightarrow \text{many-one}$$

17. If $P(X)$ represent the probability of getting a '6' in the X^{th} roll of a die for the first time. Also
 $a = P(X = 3)$
 $b = P(X \geq 3)$

$$c = P\left(\frac{X \geq 6}{X \geq 3}\right), \text{ then } \frac{b+c}{a} = ?$$

Ans. (12)

$$\text{Sol. } P(X=3) = \left(\frac{5}{6}\right)^2 \cdot \frac{1}{6} = a$$

$$P(X \geq 3) = \left(\frac{5}{6}\right)^2 = b$$

$$P\left(\frac{X \geq 6}{X > 3}\right) = \left(\frac{5}{6}\right)^2 = c$$

$$\therefore \frac{b+c}{a} = \frac{2\left(\frac{5}{6}\right)^2}{\left(\frac{5}{6}\right)^2 \cdot \frac{1}{6}} = 12$$

- 18.** If the angle between two vectors $\vec{a} = \alpha\hat{i} - 4\hat{j} - \hat{k}$ and $\vec{b} = \alpha\hat{i} + \alpha\hat{j} + 4\hat{k}$ is acute then find least positive integral value of α .

(1) 4 (2) 5

(3) 6

(4) 7

Ans. (2)

$$\text{Sol. } \vec{a} \cdot \vec{b} \geq 0$$

$$\Rightarrow a^2 - 4a - 4 \geq 0$$

$$\alpha < (2 - 2\sqrt{2}) \text{ or } \alpha > (2 + 2\sqrt{2})$$

- 19.** If $S = \{1, 2, \dots, 10\}$ and $M = P(S)$,
 If ARB such that $A \cap B \neq \emptyset$ where $A \in M, B \in M$

Then

- (1) R is reflexive and symmetric (2) Only symmetric
(3) Only reflexive (4) Symmetric and transitive

Ans. (2)

Sol. $\phi \cap \phi = \phi \Rightarrow (\phi, \phi) \notin R \Rightarrow$ not reflexive.

If $A \cap B \neq \phi \Rightarrow B \cap A \neq \phi \Rightarrow$ Symmetric

If $A \cap B \neq \phi$ and $B \cap C \neq \phi \Rightarrow A \cap C = \phi$

for example $A = \{1, 2\}$

$B = \{2, 3\}$

$C = \{3, 4\}$

- 20.** If four points $(0, 0), (1, 0), (0, 1), (2k, 3k)$ are concyclic, then k is

(1) $\frac{4}{13}$

(2) $\frac{5}{13}$

(3) $\frac{7}{13}$

(4) $\frac{9}{13}$

Ans. (2)

Sol. Equation of circle is

$$x(x-1) + y(y-1) = 0$$

$$x^2 + y^2 - x - y = 0$$

$$B(2k, 3k)$$

$$\Rightarrow 4k^2 + 9k^2 - 2k - 3k = 0$$

$$\Rightarrow 13k^2 = 5k$$

$$\Rightarrow k = 0, \frac{5}{13}$$

$$\therefore k = \frac{5}{13}$$

- 21.** If $f(x)$ is differentiable function satisfying $f(x) - f(y) \geq \log \frac{x}{y} + x - y$, then find $\sum_{N=1}^{20} f'\left(\frac{1}{N^2}\right)$

Ans. (2890)

Sol. Let $x > y$

$$\lim_{y \rightarrow x} \frac{f(x) - f(y)}{x - y} \geq \frac{\log x - \log y}{x - y} + 1$$

$$f'(x^-) \geq \frac{1}{x} + 1$$

Let $x < y$

$$\frac{f(x) - f(y)}{x - y} \leq \frac{\log x - \log y}{x - y} + 1$$

$$f'(x^+) \leq \frac{1}{x} + 1$$

$\Rightarrow f'(x^-) = f'(x^+)$ as $f(x)$ is differentiable function

$$f'(x) = \frac{1}{x} + 1$$

$$f'\left(\frac{1}{N^2}\right) = N^2 + 1$$

$$\sum_{N=1}^{20} f'\left(\frac{1}{N^2}\right) = \sum (N^2 + 1) = \frac{20 \times 21 \times 41}{6} + 20 = 2890$$

22. Let $\frac{dx}{dt} + ax = 0$ and $\frac{dy}{dt} + by = 0$ where $y(0) = 1$, $x(0) = 2$, and $x(t) = y(t)$, then t is

(1) $\frac{\ln 3}{a-b}$

(2) $\frac{\ln 2}{b-a}$

(3) $\frac{\ln 2}{a-b}$

(4) $\frac{\ln 3}{b-a}$

Ans. (3)

Sol. $\frac{dx}{dt} + ax = 0$

$$\Rightarrow \ln x = -at + c$$

$$x(0) = 2 \Rightarrow c = \ln 2$$

$$\therefore x = 2e^{-at}$$

$$\frac{dy}{dt} + by = 0 \Rightarrow y = e^{-bt}$$

$$x(t) = g(t)$$

$$2e^{-at} = e^{-bt}$$

$$\Rightarrow t = \frac{\ln 2}{a-b}$$

23. If $H(a, b)$ is the orthocentre of ΔABC where $A(1, 2)$, $B(2, 3)$ & $C(3, 1)$, then find $\frac{36I_1}{I_2}$ if

$$I_1 = \int_a^b x \sin(4x - x^2) dx \text{ and } I_2 = \int_a^b \sin(4x - x^2) dx$$

Ans. (72)

Sol. ΔABC is isosceles

$\Rightarrow H$ lies on angle bisector passing through $(3, 1)$ which is $x + y = 4$

$$\therefore a + b = 4$$

Now apply $(a + b - x)$ in I_1

$$2I_1 = \int_a^b 4 \sin(4x - x^2) dx$$

$$\Rightarrow 2I_1 = 4I_2$$

$$\Rightarrow \frac{I_1}{I_2} = 2$$

$$\therefore \frac{36I_1}{I_2} = 72$$

24. $f(x) = \begin{cases} \frac{\sin(x-3)}{2^{x-[x]}} & , \quad x > 3 \\ -\frac{a(x^2 - 7x + 12)}{b|x^2 - 7x + 12|} & , \quad x < 3 \\ b & , \quad x = 3 \end{cases}$. Find number of ordered pairs (a, b) so that f(x) is continuous at x = 3

Ans. (1)
Sol. LHL = RHL = f(3)

$$-\frac{a}{b} = 2^1 = b$$

$$\Rightarrow b = 2 \text{ and } a = -4$$

$$\Rightarrow (a, b) = (-4, 2)$$

25. Let $A = \begin{bmatrix} 2 & 0 & 1 \\ 1 & 0 & 0 \\ 3 & 2 & 0 \end{bmatrix}$, $B = [B_1 \ B_2 \ B_3]$ where B_1, B_2, B_3 are column matrices such that

$$AB_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, AB_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, AB_3 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

α = sum of diagonal elements of B

$$\beta = |B|, \text{ then find } |\alpha^3 + \beta^3|$$

Ans. (1.125)

Sol. $A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & -\frac{3}{2} & \frac{1}{2} \\ 1 & -2 & 0 \end{bmatrix}$

$$B_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, B_2 = \begin{bmatrix} 0 \\ \frac{1}{2} \\ 2 \end{bmatrix}, B_3 = \begin{bmatrix} 2 \\ -\frac{5}{2} \\ -1 \end{bmatrix}$$

$$\text{Tr}(B) = -\frac{1}{2}$$

$$|B| = -1$$

$$\therefore a = -\frac{1}{2}, b = -1$$

$$|\alpha^3 + \beta^3| = \frac{9}{8} = 1.125$$

26. If $\cos(2x) - a \sin x = 2a - 7$ has a solution for $a \in [p, q]$ and $r = \tan 9^\circ + \tan 63^\circ + \tan 81^\circ + \tan 27^\circ$, then $p \cdot q \cdot r = ?$

(1) $40\sqrt{5}$ (2) $32\sqrt{5}$ (3) $30\sqrt{5}$ (4) $48\sqrt{5}$

Ans. (4)

Sol. $2(\sin^2 x - 4) + a(\sin x + 2) = 0$

$$2(\sin x - 2) + a = 0$$

$$\Rightarrow a = 4 - 2 \sin x$$

$$a \in [2, 6]$$

$$\text{Also, } r = \left(\tan 9^\circ + \frac{1}{\tan 9^\circ} \right) + \left(\tan 27^\circ + \frac{1}{\tan 27^\circ} \right)$$

$$= \frac{2}{\sin 18^\circ} + \frac{2}{\sin 54^\circ}$$

$$= \frac{2 \times 4}{\sqrt{5}-1} + \frac{2 \times 4}{\sqrt{5}+1}$$

$$= \frac{8 \times 2\sqrt{5}}{4} = 4\sqrt{5}$$

$$\therefore pqr = 48\sqrt{5}$$



RANKERS